

## Cold-induced sweating syndrome

### Description

Cold-induced sweating syndrome is characterized by problems with regulating body temperature and other abnormalities affecting many parts of the body. In infancy, the features of this condition are often known as Crisponi syndrome. Researchers originally thought that cold-induced sweating syndrome and Crisponi syndrome were separate disorders, but it is now widely believed that they represent the same condition at different times during life.

Infants with Crisponi syndrome have unusual facial features, including a flat nasal bridge, upturned nostrils, a long space between the nose and upper lip (philtrum), a high arched roof of the mouth (palate), a small chin (micrognathia), and low-set ears. The muscles in the lower part of the face are weak, leading to severe feeding difficulties, excessive drooling, and breathing problems. Other physical abnormalities associated with Crisponi syndrome include a scaly skin rash, an inability to fully extend the elbows, overlapping fingers and tightly fisted hands, and malformations of the feet and toes. Affected infants startle easily and often tense their facial muscles into a grimace-like expression. By six months of age, infants with Crisponi syndrome develop unexplained high fevers that increase the risk of seizures and sudden death.

Many of the health problems associated with Crisponi syndrome improve with time, and affected individuals who survive the newborn period go on to develop other features of cold-induced sweating syndrome in early childhood. Within the first decade of life, affected individuals begin having episodes of profuse sweating (hyperhidrosis) and shivering involving the face, torso, and arms. The excessive sweating is usually triggered by exposure to temperatures below about 65 or 70 degrees Fahrenheit, but it can also be triggered by nervousness or eating sugary foods. Paradoxically, affected individuals tend not to sweat in warmer conditions, instead becoming flushed and overheated in hot environments.

Adolescents with cold-induced sweating syndrome typically develop abnormal side-to-side and front-to-back curvature of the spine (scoliosis and kyphosis, often called kyphoscoliosis when they occur together). Although infants may develop life-threatening fevers, affected individuals who survive infancy have a normal life expectancy.

## Frequency

Cold-induced sweating syndrome is a rare condition; its prevalence is unknown. The condition was first identified in the Sardinian population, but it has since been reported in regions worldwide.

## Causes

About 90 percent of cases of cold-induced sweating syndrome and Crisponi syndrome result from mutations in the *CRLF1* gene. These cases are designated as CISS1. The remaining 10 percent of cases are caused by mutations in the *CLCF1* gene and are designated as CISS2.

The proteins produced from the *CRLF1* and *CLCF1* genes work together as part of a signaling pathway that is involved in the normal development of the nervous system. This pathway appears to be particularly important for the development and maintenance of motor neurons, which are nerve cells that control muscle movement. Studies suggest that this pathway also has a role in a part of the nervous system known as the sympathetic nervous system, specifically in the regulation of sweating in response to temperature changes and other factors. The proteins produced from the *CRLF1* and *CLCF1* genes appear to be critical for the normal development and maturation of nerve cells that control the activity of sweat glands.

Additionally, the *CRLF1* and *CLCF1* genes likely have functions outside the nervous system, including roles in the body's inflammatory response and in bone development. However, little is known about their involvement in these processes.

Mutations in either the *CRLF1* or *CLCF1* gene disrupt the normal development of several body systems, including the nervous system. The role of these genes in sympathetic nervous system development may help explain the abnormal sweating that is characteristic of this condition, including unusual sweating patterns and related problems with body temperature regulation. The involvement of these genes in motor neuron development and bone development provides clues to some of the other signs and symptoms of cold-induced sweating syndrome, including distinctive facial features, facial muscle weakness, and skeletal abnormalities. However, little is known about how *CRLF1* or *CLCF1* gene mutations underlie these other features of cold-induced sweating syndrome.

[Learn more about the genes associated with Cold-induced sweating syndrome](#)

- *CLCF1*
- *CRLF1*

## Inheritance

Cold-induced sweating syndrome is inherited in an autosomal recessive pattern, which means both copies of the *CRLF1* or *CLCF1* gene in each cell have mutations. The

parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## **Other Names for This Condition**

- CISS
- CNTF receptor-related disorders
- Crisponi syndrome
- Sohar-Crisponi syndrome

## **Additional Information & Resources**

### Genetic Testing Information

- Genetic Testing Registry: Cold-induced sweating syndrome (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1832409/>)
- Genetic Testing Registry: Cold-induced sweating syndrome 1 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1848947/>)
- Genetic Testing Registry: Cold-induced sweating syndrome 2 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1853198/>)
- Genetic Testing Registry: PERCHING syndrome (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C4310742/>)

### Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

### Catalog of Genes and Diseases from OMIM

- CRISPONI/COLD-INDUCED SWEATING SYNDROME 1; CISS1 (<https://omim.org/entry/272430>)
- CRISPONI/COLD-INDUCED SWEATING SYNDROME 2; CISS2 (<https://omim.org/entry/610313>)
- PERCHING SYNDROME; PERCHING (<https://omim.org/entry/617055>)

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28cold-induced+sweating+syndrome%5BTIAB%5D%29+OR+%28crisponi+syndrome%5BTIAB%5D%29+OR+%28CISS1%5BTIAB%5D%29+OR+%28CISS2%5BTIAB%5D%29%29+AND+englis h%5Bla%5D+AND+human%5Bmh%5D>)

## References

- Angius A, Uva P, Buers I, Oppo M, Puddu A, Onano S, Persico I, Loi A, Marcia L, Hohne W, Cuccuru G, Fotia G, Deiana M, Marongiu M, Atalay HT, Inan S, El Assy O, Smit LM, Okur I, Boduroglu K, Utine GE, Kilic E, Zampino G, Crisponi G, Crisponi L, Rutsch F. Bi-allelic Mutations in KLHL7 Cause a Crisponi/CISS1-like Phenotype Associated with Early-Onset Retinitis Pigmentosa. *Am J Hum Genet.* 2016 Jul 7;99(1):236-45. doi: 10.1016/j.ajhg.2016.05.026. Erratum In: *Am J Hum Genet.* 2018 Apr 5;102(4):713. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27392078>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5005468/>)
- Crisponi G. Autosomal recessive disorder with muscle contractions resembling neonatal tetanus, characteristic face, camptodactyly, hyperthermia, and sudden death: a new syndrome? *Am J Med Genet.* 1996 Apr 24;62(4):365-71. doi:10.1002/(SICI)1096-8628(19960424)62:43.0.CO;2-Q. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8723066>)
- Crisponi L, Crisponi G, Meloni A, Toliat MR, Nurnberg G, Usala G, Uda M, Masala M, Hohne W, Becker C, Marongiu M, Chiappe F, Kleta R, Rauch A, Wollnik B, Strasser F, Reese T, Jakobs C, Kurlermann G, Cao A, Nurnberg P, Rutsch F. Crisponi syndrome is caused by mutations in the CRLF1 gene and is allelic to cold-induced sweating syndrome type 1. *Am J Hum Genet.* 2007 May;80(5):971-81. doi: 10.1086/516843. Epub 2007 Mar 30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17436252>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1852730/>)
- Dagoneau N, Bellais S, Blanchet P, Sarda P, Al-Gazali LI, Di Rocco M, Huber C, Djouadi F, Le Goff C, Munnich A, Cormier-Daire V. Mutations in cytokine receptor-like factor 1 (CRLF1) account for both Crisponi and cold-induced sweating syndromes. *Am J Hum Genet.* 2007 May;80(5):966-70. doi: 10.1086/513608. Epub 2007 Mar 13. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17436251>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1852726/>)
- Hahn AF, Jones DL, Knappskog PM, Boman H, McLeod JG. Cold-induced sweating syndrome: a report of two cases and demonstration of genetic heterogeneity. *J Neurol Sci.* 2006 Dec 1;250(1-2):62-70. doi: 10.1016/j.jns.2006.07.001. Epub 2006 Sep 6. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16952376>)
- Hahn AF, Knappskog PM. Cold-Induced Sweating Syndrome Including Crisponi Syndrome. 2011 Mar 3 [updated 2021 Aug 12]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews*(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK52917/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21370513>)
- Hahn AF, Waaler PE, Kvistad PH, Bamforth JS, Miles JH, McLeod JG, Knappskog PM, Boman H. Cold-induced sweating syndrome: CISS1 and CISS2: manifestations from infancy to adulthood. Four new cases. *J Neurol Sci.* 2010 Jun 15;293(1-2):68-75. doi: 10.1016/j.jns.2010.02.028. Epub 2010 Apr 18. Citation on

PubMed (<https://pubmed.ncbi.nlm.nih.gov/20400119>)

- Knappskog PM, Majewski J, Livneh A, Nilsen PT, Bringsli JS, Ott J, Boman H. Cold-induced sweating syndrome is caused by mutations in the CRLF1 gene. *Am J Hum Genet.* 2003 Feb;72(2):375-83. doi: 10.1086/346120. Epub 2002 Dec 31. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12509788>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC379230/>)
- Rousseau F, Gauchat JF, McLeod JG, Chevalier S, Guillet C, Guilhot F, Cognetl, Froger J, Hahn AF, Knappskog PM, Gascan H, Boman H. Inactivation of cardiotrophin-like cytokine, a second ligand for ciliary neurotrophic factor receptor, leads to cold-induced sweating syndrome in a patient. *Proc Natl Acad Sci U S A.* 2006 Jun 27;103(26):10068-73. doi: 10.1073/pnas.0509598103. Epub 2006 Jun 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16782820>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1502507/>)

**Last updated August 1, 2012**