

Combined pituitary hormone deficiency

Description

Combined pituitary hormone deficiency is a condition that causes a shortage (deficiency) of several hormones produced by the pituitary gland, which is located at the base of the brain. A lack of these hormones may affect the development of many parts of the body. The first signs of this condition include a failure to grow at the expected rate and short stature that usually becomes apparent in early childhood.

People with combined pituitary hormone deficiency may have hypothyroidism, which is underactivity of the butterfly-shaped thyroid gland in the lower neck. Hypothyroidism can cause many symptoms, including weight gain and fatigue. Other features of combined pituitary hormone deficiency include delayed or absent puberty and lack the ability to have biological children (infertility). The condition can also be associated with a deficiency of the hormone cortisol. Cortisol deficiency can impair the body's immune system, causing individuals to be more susceptible to infection.

Rarely, people with combined pituitary hormone deficiency have intellectual disability; a short, stiff neck; or underdeveloped optic nerves, which carry visual information from the eyes to the brain.

Frequency

The prevalence of combined pituitary hormone deficiency is estimated to be 1 in 8,000 individuals worldwide.

Causes

Mutations in at least eight genes have been found to cause combined pituitary hormone deficiency. Mutations in the *PROP1* gene are the most common known cause of this disorder, accounting for an estimated 12 to 55 percent of cases. Mutations in other genes have each been identified in a smaller number of affected individuals.

The genes associated with combined pituitary hormone deficiency provide instructions for making proteins called transcription factors, which help control the activity of many other genes. The proteins are involved in the development of the pituitary gland and the specialization (differentiation) of its cell types. The cells of the pituitary gland are responsible for triggering the release of several hormones that direct the development of many parts of the body. Some of the transcription factors are found only in the

pituitary gland, and some are also active in other parts of the body.

Mutations in the genes associated with combined pituitary hormone deficiency can result in abnormal differentiation of pituitary gland cells and may prevent the production of several hormones. These hormones can include growth hormone (GH), which is needed for normal growth; follicle-stimulating hormone (FSH) and luteinizing hormone (LH), which both play a role in sexual development and the ability to have children (fertility); thyroid-stimulating hormone (TSH), which helps with thyroid gland function; prolactin, which stimulates the production of breast milk; and adrenocorticotrophic hormone (ACTH), which influences energy production in the body and maintains normal blood glucose and blood pressure levels. The degree to which these hormones are deficient is variable, with prolactin and ACTH showing the most variability. In many affected individuals, ACTH deficiency does not occur until late adulthood.

Most people with combined pituitary hormone deficiency do not have identified mutations in any of the genes known to be associated with this condition. The cause of the disorder in these individuals is unknown.

[Learn more about the genes associated with Combined pituitary hormone deficiency](#)

- HESX1
- OTX2
- PROKR2
- PROP1
- SOX2

Additional Information from NCBI Gene:

- GLI2
- LHX3
- LHX4
- POU1F1

Inheritance

Most cases of combined pituitary hormone deficiency are sporadic, which means they occur in people with no history of the disorder in their family. Less commonly, this condition has been found to run in families. When the disorder is familial, it can have an autosomal dominant or an autosomal recessive pattern of inheritance.

Autosomal dominant inheritance means one copy of an altered gene in each cell is sufficient to cause the disorder.

Autosomal recessive inheritance means both copies of a gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each

carry one copy of a mutated gene, but they typically do not show signs and symptoms of the condition. Most cases of familial combined pituitary hormone deficiency are inherited in an autosomal recessive pattern.

Other Names for This Condition

- CPHD
- Panhypopituitarism

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Combined pituitary hormone deficiencies, genetic form (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C4273747/>)

Genetic and Rare Diseases Information Center

- Combined pituitary hormone deficiencies, genetic forms (<https://rarediseases.info.nih.gov/diseases/10602/index>)
- Non-acquired combined pituitary hormone deficiency-sensorineural hearing loss-spine abnormalities syndrome (<https://rarediseases.info.nih.gov/diseases/10603/index>)
- Pituitary hormone deficiency, combined 1 (<https://rarediseases.info.nih.gov/diseases/10601/index>)
- Short stature-pituitary and cerebellar defects-small sella turcica syndrome (<https://rarediseases.info.nih.gov/diseases/10604/index>)

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Clinical Trials

- ClinicalTrials.gov ([https://clinicaltrials.gov/search?cond=%22Combined pituitary hormone deficiency%22](https://clinicaltrials.gov/search?cond=%22Combined%20pituitary%20hormone%20deficiency%22))

Catalog of Genes and Diseases from OMIM

- PITUITARY HORMONE DEFICIENCY, COMBINED, 3; CPHD3 (<https://omim.org/entry/221750>)
- PITUITARY HORMONE DEFICIENCY, COMBINED, 2; CPHD2 (<https://omim.org/entry/262600>)

- PITUITARY HORMONE DEFICIENCY, COMBINED, 4; CPHD4 (<https://omim.org/entry/262700>)
- PITUITARY HORMONE DEFICIENCY, COMBINED OR ISOLATED, 1; CPHD1 (<https://omim.org/entry/613038>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28combined+pituitary+hormone+deficiency%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>)

References

- Kelberman D, Rizzoti K, Lovell-Badge R, Robinson IC, Dattani MT. Genetic regulation of pituitary gland development in human and mouse. *Endocr Rev.* 2009Dec;30(7):790-829. doi: 10.1210/er.2009-0008. Epub 2009 Oct 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19837867>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2806371/>)
- Kelberman D, Turton JP, Woods KS, Mehta A, Al-Khawari M, Greening J, Swift PG, Otonkoski T, Rhodes SJ, Dattani MT. Molecular analysis of novel PROP1 mutations associated with combined pituitary hormone deficiency (CPHD). *Clin Endocrinol(Oxf).* 2009 Jan;70(1):96-103. doi: 10.1111/j.1365-2265.2008.03326.x. Epub 2008 Jun 25. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19128366>)
- Mody S, Brown MR, Parks JS. The spectrum of hypopituitarism caused by PROP1 mutations. *Best Pract Res Clin Endocrinol Metab.* 2002 Sep;16(3):421-31. doi:10.1053/beem.2002.0218. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12464226>)
- Reynaud R, Gueydan M, Saveanu A, Vallette-Kasic S, Enjalbert A, Brue T, Barlier A. Genetic screening of combined pituitary hormone deficiency: experience in 195 patients. *J Clin Endocrinol Metab.* 2006 Sep;91(9):3329-36. doi:10.1210/jc.2005-2173. Epub 2006 May 30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16735499>)
- Vieira TC, Boldarine VT, Abucham J. Molecular analysis of PROP1, PIT1, HESX1, LHX3, and LHX4 shows high frequency of PROP1 mutations in patients with familial forms of combined pituitary hormone deficiency. *Arq Bras Endocrinol Metabol.* 2007 Oct;51(7):1097-103. doi: 10.1590/s0004-27302007000700012. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18157385>)

Last updated August 1, 2010