

Epidermolytic hyperkeratosis

Description

Epidermolytic hyperkeratosis is a skin disorder that is present at birth. Affected babies may have very red skin (erythroderma) and severe blisters. Because newborns with this disorder are missing the protection provided by normal skin, they are at risk of becoming dehydrated and developing infections in the skin or throughout the body (sepsis).

As affected individuals get older, blistering is less frequent, erythroderma becomes less evident, and the skin becomes thick (hyperkeratotic), especially over joints, on areas of skin that come into contact with each other, or on the scalp or neck. This thickened skin is usually darker than normal. Bacteria can grow in the thick skin, often causing a distinct odor.

Epidermolytic hyperkeratosis can be categorized into two types. People with PS-type epidermolytic hyperkeratosis have thick skin on the palms of their hands and soles of their feet (palmoplantar or palm/sole hyperkeratosis) in addition to other areas of the body. People with the other type, NPS-type, do not have extensive palmoplantar hyperkeratosis but do have hyperkeratosis on other areas of the body.

Epidermolytic hyperkeratosis is part of a group of conditions called ichthyoses, which refers to the scaly skin seen in individuals with related disorders. However, in epidermolytic hyperkeratosis, the skin is thick but not scaly as in some of the other conditions in the group.

Frequency

Epidermolytic hyperkeratosis affects approximately 1 in 200,000 to 300,000 people worldwide.

Causes

Mutations in the *KRT1* or *KRT10* genes are responsible for epidermolytic hyperkeratosis. These genes provide instructions for making proteins called keratin 1 and keratin 10, which are found in cells called keratinocytes in the outer layer of the skin (the epidermis). The tough, fibrous keratin proteins attach to each other and form fibers called intermediate filaments, which form networks and provide strength and resiliency to the epidermis.

Mutations in the *KRT1* or *KRT10* genes lead to changes in the keratin proteins, preventing them from forming strong, stable intermediate filament networks within cells. Without a strong network, keratinocytes become fragile and are easily damaged, which can lead to blistering in response to friction or mild trauma. It is unclear how these mutations cause the overgrowth of epidermal cells that results in hyperkeratotic skin.

KRT1 gene mutations are associated with PS-type epidermal hyperkeratosis, and *KRT10* gene mutations are usually associated with NPS-type. The keratin 1 protein is present in the keratinocytes of the skin on the palms of the hands and the soles of the feet as well as other parts of the body, so mutations in the *KRT1* gene lead to skin problems in these areas. The keratin 10 protein is not found in the skin of the palms and soles, so these areas are unaffected by mutations in the *KRT10* gene.

[Learn more about the genes associated with Epidermolytic hyperkeratosis](#)

- KRT1
- KRT10

Inheritance

Epidermolytic hyperkeratosis can have different inheritance patterns. About half of the cases of this condition result from new mutations in the *KRT1* or *KRT10* gene and occur in people with no history of the disorder in their family.

When epidermolytic hyperkeratosis is inherited, it is usually in an autosomal dominant pattern, which means one copy of the altered *KRT1* or *KRT10* gene in each cell is sufficient to cause the disorder.

Very rarely, epidermolytic hyperkeratosis caused by mutations in the *KRT10* gene can be inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

Other Names for This Condition

- BCIE
- BIE
- Bullous congenital ichthyosiform erythroderma
- Bullous erythroderma ichthyosiforme
- Bullous erythroderma ichthyosiformis congenita of Brocq
- Bullous ichthyosiform erythroderma
- EHK
- Epidermolytic ichthyosis
- Hyperkeratosis, epidermolytic

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Bullous ichthyosiform erythroderma (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0079153/>)

Genetic and Rare Diseases Information Center

- Autosomal dominant epidermolytic ichthyosis (<https://rarediseases.info.nih.gov/diseases/1039/index>)

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Clinical Trials

- ClinicalTrials.gov ([https://clinicaltrials.gov/search?cond=%22Epidermolytic hyperkeratosis%22](https://clinicaltrials.gov/search?cond=%22Epidermolytic%20hyperkeratosis%22))

Catalog of Genes and Diseases from OMIM

- EPIDERMOLYTIC HYPERKERATOSIS 1; EHK1 (<https://omim.org/entry/113800>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28Hyperkeratosis,+Epidermolytic%5BMAJR%5D%29+AND+%28epidermolytic+hyperkeratosis%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

References

- Chamcheu JC, Siddiqui IA, Syed DN, Adhami VM, Liovic M, Mukhtar H. Keratogene mutations in disorders of human skin and its appendages. Arch Biochem Biophys. 2011 Apr 15;508(2):123-37. doi: 10.1016/j.abb.2010.12.019. Epub 2010 Dec 19. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21176769>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3142884/>)
- Chipev CC, Korge BP, Markova N, Bale SJ, DiGiovanna JJ, Compton JG, Steinert PM. A leucine----proline mutation in the H1 subdomain of keratin 1 causes epidermolytic hyperkeratosis. Cell. 1992 Sep 4;70(5):821-8. doi:10.1016/

0092-8674(92)90315-4. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/1381288>)

- DiGiovanna JJ, Bale SJ. Clinical heterogeneity in epidermolytic hyperkeratosis. *Arch Dermatol*. 1994 Aug;130(8):1026-35. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8053700>)
- Yang JM, Chipev CC, DiGiovanna JJ, Bale SJ, Marekov LN, Steinert PM, Compton JG. Mutations in the H1 and 1A domains in the keratin 1 gene in epidermolytic hyperkeratosis. *J Invest Dermatol*. 1994 Jan;102(1):17-23. doi:10.1111/1523-1747.ep12371725. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/7507151>)
- Yang JM, Nam K, Kim HC, Lee JH, Park JK, Wu K, Lee ES, Steinert PM. A novel glutamic acid to aspartic acid mutation near the end of the 2B rod domain in the keratin 1 chain in epidermolytic hyperkeratosis. *J Invest Dermatol*. 1999 Mar;112(3):376-9. doi: 10.1038/sj.jid.5600439. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10084317>)

Last updated November 1, 2011