

Lattice corneal dystrophy type I

Description

Lattice corneal dystrophy type I is an eye disorder that affects the clear, outer covering of the eye called the cornea. The cornea must remain clear for an individual to see properly; however, in lattice corneal dystrophy type I, protein clumps known as amyloid deposits cloud the cornea, which leads to vision impairment. The cornea is made up of several layers of tissue, and in lattice corneal dystrophy type I, the deposits form in the stromal layer. The amyloid deposits form as delicate, branching fibers that create a lattice pattern.

Affected individuals often have recurrent corneal erosions, which are caused by separation of particular layers of the cornea from one another. Corneal erosions are very painful and can cause sensitivity to bright light (photophobia). Lattice corneal dystrophy type I is usually bilateral, which means it affects both eyes. The condition becomes apparent in childhood or adolescence and leads to vision problems by early adulthood.

Frequency

Lattice corneal dystrophy type I is one of the most common disorders in a group of conditions that are characterized by protein deposits in the cornea (corneal dystrophies); however, it is still a rare condition. The prevalence of lattice corneal dystrophy type I is unknown.

Causes

Lattice corneal dystrophy type I is caused by mutations in the *TGFBI* gene. This gene provides instructions for making a protein that is found in many tissues throughout the body, including the cornea. The TGFBI protein is part of the extracellular matrix, an intricate network that forms in the spaces between cells and provides structural support to tissues. The protein is thought to play a role in the attachment of cells to one another (cell adhesion) and cell movement (migration).

The *TGFBI* gene mutations involved in lattice corneal dystrophy type I change single protein building blocks (amino acids) in the TGFBI protein. Mutated TGFBI proteins abnormally clump together and form amyloid deposits. However, it is unclear how the changes caused by the gene mutations induce the protein to form deposits.

[Learn more about the gene associated with Lattice corneal dystrophy type I](#)

- TGFB1

Inheritance

This condition is inherited in an autosomal dominant pattern, which means one copy of the altered gene in each cell is sufficient to cause the disorder. In most cases, an affected person has one parent with the condition.

Other Names for This Condition

- Biber-Haab-Dimmer dystrophy

Additional Information & Resources

Genetic Testing Information

- Genetic Testing Registry: Lattice corneal dystrophy Type I (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1690006/>)

Genetic and Rare Diseases Information Center

- Lattice corneal dystrophy type I (<https://rarediseases.info.nih.gov/diseases/9678/index>)

Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

Catalog of Genes and Diseases from OMIM

- CORNEAL DYSTROPHY, LATTICE TYPE I; CDL1 (<https://omim.org/entry/122200>)

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28lattice+corneal+dystrophy+type+i%5BTIAB%5D%29+OR+%28lattice+corneal+dystrophy+type+1%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>)

References

- Kannabiran C, Klintworth GK. TGFBI gene mutations in corneal dystrophies. *HumMutat.* 2006 Jul;27(7):615-25. doi: 10.1002/humu.20334. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16683255>)
- Klintworth GK. Corneal dystrophies. *Orphanet J Rare Dis.* 2009 Feb 23;4:7. doi:10.1186/1750-1172-4-7. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19236704>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2695576/>)
- Liu Z, Wang YQ, Gong QH, Xie LX. An R124C mutation in TGFBI caused latticecorneal dystrophy type I with a variable phenotype in three Chinese families. *MolVis.* 2008 Jun 30;14:1234-9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18615206>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2443752/>)
- Munier FL, Korvatska E, Djemai A, Le Paslier D, Zografos L, Pescia G, Schorderet DF. Kerato-epithelin mutations in four 5q31-linked cornealdystrophies. *Nat Genet.* 1997 Mar;15(3):247-51. doi: 10.1038/ng0397-247. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9054935>)
- Schmitt-Bernard CF, Chavanieu A, Derancourt J, Arnaud B, Demaille JG, Calas B, Argiles A. In vitro creation of amyloid fibrils from native and Arg124Cys mutatedbetalGH3((110-131)) peptides, and its relevance for lattice corneal amyloiddystrophy type I. *Biochem Biophys Res Commun.* 2000 Jul 5;273(2):649-53. doi:10.1006/bbrc.2000.2955. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10873659>)

Last updated April 1, 2012