

## Primary ciliary dyskinesia

### Description

Primary ciliary dyskinesia is a disorder characterized by chronic respiratory tract infections, abnormally positioned internal organs, and the inability to have children (infertility). The signs and symptoms of this condition are caused by abnormal cilia and flagella. Cilia are microscopic, finger-like projections that stick out from the surface of cells. They are found in the linings of the airway, the reproductive system, and other organs and tissues. Flagella are tail-like structures, similar to cilia, that propel sperm cells forward.

In the respiratory tract, cilia move back and forth in a coordinated way to move mucus towards the throat. This movement of mucus helps to eliminate fluid, bacteria, and particles from the lungs. Most babies with primary ciliary dyskinesia experience breathing problems at birth, which suggests that cilia play an important role in clearing fetal fluid from the lungs. Beginning in early childhood, affected individuals develop frequent respiratory tract infections. Without properly functioning cilia in the airway, bacteria remain in the respiratory tract and cause infection. People with primary ciliary dyskinesia also have year-round nasal congestion and a chronic cough. Chronic respiratory tract infections can result in a condition called bronchiectasis, which damages the passages, called bronchi, leading from the windpipe to the lungs and can cause life-threatening breathing problems.

Some individuals with primary ciliary dyskinesia have abnormally placed organs within their chest and abdomen. These abnormalities arise early in embryonic development when the differences between the left and right sides of the body are established. About 50 percent of people with primary ciliary dyskinesia have a mirror-image reversal of their internal organs (situs inversus totalis). For example, in these individuals the heart is on the right side of the body instead of on the left. Situs inversus totalis does not cause any apparent health problems. When someone with primary ciliary dyskinesia has situs inversus totalis, they are often said to have Kartagener syndrome.

Approximately 12 percent of people with primary ciliary dyskinesia have a condition known as heterotaxy syndrome or situs ambiguus, which is characterized by abnormalities of the heart, liver, intestines, or spleen. These organs may be structurally abnormal or improperly positioned. In addition, affected individuals may lack a spleen (asplenia) or have multiple spleens (polysplenia). Heterotaxy syndrome results from problems establishing the left and right sides of the body during embryonic development. The severity of heterotaxy varies widely among affected individuals.

Primary ciliary dyskinesia can also lead to infertility. Vigorous movements of the flagella are necessary to propel the sperm cells forward to the female egg cell. Because their sperm do not move properly, males with primary ciliary dyskinesia are usually unable to father children. Infertility occurs in some affected females and is likely due to abnormal cilia in the fallopian tubes.

Another feature of primary ciliary dyskinesia is recurrent ear infections (otitis media), especially in young children. Otitis media can lead to permanent hearing loss if untreated. The ear infections are likely related to abnormal cilia within the inner ear.

Rarely, individuals with primary ciliary dyskinesia have an accumulation of fluid in the brain (hydrocephalus), likely due to abnormal cilia in the brain.

## Frequency

Primary ciliary dyskinesia occurs in approximately 1 in 16,000 individuals.

## Causes

Primary ciliary dyskinesia can result from mutations in many different genes. These genes provide instructions for making proteins that form the inner structure of cilia and produce the force needed for cilia to bend. Coordinated back and forth movement of cilia is necessary for the normal functioning of many organs and tissues. The movement of cilia also helps establish the left-right axis (the imaginary line that separates the left and right sides of the body) during embryonic development.

Mutations in the genes that cause primary ciliary dyskinesia result in defective cilia that move abnormally or are unable to move (immotile). Because cilia have many important functions within the body, defects in these cell structures cause a variety of signs and symptoms.

Mutations in the *DNAI1* and *DNAH5* genes account for up to 30 percent of all cases of primary ciliary dyskinesia. Mutations in the other genes associated with this condition are found in only a small percentage of cases. In many people with primary ciliary dyskinesia, the cause of the disorder is unknown.

[Learn more about the genes associated with Primary ciliary dyskinesia](#)

- DNAH5
- DNAI1
- OFD1
- RPGR

## Additional Information from NCBI Gene:

- CCDC103
- CCDC39

- CCDC40
- CCDC65
- CFAP298
- DNAAF1
- DNAAF11
- DNAAF2
- DNAAF3
- DNAAF4
- DNAAF5
- DNAH11
- DNAH8
- DNAI2
- DNAL1
- DRC1
- HYDIN
- NME8
- ODAD1
- ODAD2
- RSPH1
- RSPH4A
- RSPH9
- SPAG1
- ZMYND10

## **Inheritance**

This condition is inherited in an autosomal recessive pattern, which means both copies of the gene in each cell have mutations. The parents of an individual with an autosomal recessive condition each carry one copy of the mutated gene, but they typically do not show signs and symptoms of the condition.

## **Other Names for This Condition**

- Immotile cilia syndrome
- PCD

## **Additional Information & Resources**

### Genetic Testing Information

- Genetic Testing Registry: Primary ciliary dyskinesia (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C0008780/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 10 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2675867/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 11 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2675229/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 12 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2675228/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 13 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2750790/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 14 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3151136/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 15 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3151137/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 16 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3151460/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 17 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C3542550/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 2 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1847554/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 3 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1837618/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 4 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1837616/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 5 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1837615/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 6 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C1970506/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 7 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2678473/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 8 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2677085/>)
- Genetic Testing Registry: Primary ciliary dyskinesia 9 (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C2676235/>)
- Genetic Testing Registry: Kartagener syndrome (<https://www.ncbi.nlm.nih.gov/gtr/conditions/C4551906/>)

### Genetic and Rare Diseases Information Center

- Kartagener syndrome (<https://rarediseases.info.nih.gov/diseases/6815/index>)
- Primary ciliary dyskinesia (<https://rarediseases.info.nih.gov/diseases/4484/index>)

### Patient Support and Advocacy Resources

- National Organization for Rare Disorders (NORD) (<https://rarediseases.org/>)

### Clinical Trials

- ClinicalTrials.gov ([https://clinicaltrials.gov/search?cond=%22Primary ciliary dyskinesia%22](https://clinicaltrials.gov/search?cond=%22Primary+ciliary+dyskinesia%22))

### Catalog of Genes and Diseases from OMIM

- CILIARY DYSKINESIA, PRIMARY, 1; CILD1 (<https://omim.org/entry/244400>)
- CILIARY DYSKINESIA, PRIMARY, 3; CILD3 (<https://omim.org/entry/608644>)
- CILIARY DYSKINESIA, PRIMARY, 4; CILD4 (<https://omim.org/entry/608646>)
- CILIARY DYSKINESIA, PRIMARY, 5; CILD5 (<https://omim.org/entry/608647>)
- CILIARY DYSKINESIA, PRIMARY, 2; CILD2 (<https://omim.org/entry/606763>)
- CILIARY DYSKINESIA, PRIMARY, 6; CILD6 (<https://omim.org/entry/610852>)
- CILIARY DYSKINESIA, PRIMARY, 8; CILD8 (<https://omim.org/entry/612274>)
- CILIARY DYSKINESIA, PRIMARY, 10; CILD10 (<https://omim.org/entry/612518>)
- CILIARY DYSKINESIA, PRIMARY, 7; CILD7 (<https://omim.org/entry/611884>)
- CILIARY DYSKINESIA, PRIMARY, 9; CILD9 (<https://omim.org/entry/612444>)
- CILIARY DYSKINESIA, PRIMARY, 13; CILD13 (<https://omim.org/entry/613193>)
- CILIARY DYSKINESIA, PRIMARY, 14; CILD14 (<https://omim.org/entry/613807>)
- CILIARY DYSKINESIA, PRIMARY, 15; CILD15 (<https://omim.org/entry/613808>)
- CILIARY DYSKINESIA, PRIMARY, 11; CILD11 (<https://omim.org/entry/612649>)
- CILIARY DYSKINESIA, PRIMARY, 12; CILD12 (<https://omim.org/entry/612650>)

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28primary+ciliary+dyskinesia%5BTI%5D%29+OR+%28immotile+cilia+syndrome%5BTI%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>)

## References

- Escudier E, Duquesnoy P, Papon JF, Amselem S. Ciliary defects and genetics of primary ciliary dyskinesia. *Paediatr Respir Rev*. 2009 Jun;10(2):51-4. doi:10.1016/j.prrv.2009.02.001. Epub 2009 Apr 18. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19410201>)
- Faily M, Bartoloni L, Letourneau A, Munoz A, Falconnet E, Rossier C, de SantiMM, Santamaria F, Sacco O, DeLozier-Blanchet CD, Lazor R, Blouin JL. Mutations in DNAH5 account for only 15% of a non-preselected cohort of patients with primary ciliary dyskinesia. *J Med Genet*. 2009 Apr;46(4):281-6. doi:10.1136/jmg.2008.061176. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19357118>)
- Faily M, Saitta A, Munoz A, Falconnet E, Rossier C, Santamaria F, de SantiMM, Lazor R, DeLozier-Blanchet CD, Bartoloni L, Blouin JL. DNAI1 mutations explain only 2% of primary ciliary dyskinesia. *Respiration*. 2008;76(2):198-204. doi: 10.1159/000128567. Epub 2008 Apr 23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18434704>)
- Horani A, Brody SL, Ferkol TW. Picking up speed: advances in the genetics of primary ciliary dyskinesia. *Pediatr Res*. 2014 Jan;75(1-2):158-64. doi:10.1038/pr.2013.200. Epub 2013 Nov 5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24192704>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3946436/>)
- Hornef N, Olbrich H, Horvath J, Zariwala MA, Fliegau M, Loges NT, Wildhaber J, Noone PG, Kennedy M, Antonarakis SE, Blouin JL, Bartoloni L, Nusslein T, Ahrens P, Griesse M, Kuhl H, Sudbrak R, Knowles MR, Reinhardt R, Omran H. DNAH5 mutations are a common cause of primary ciliary dyskinesia with outer dynein arm defects. *Am J Respir Crit Care Med*. 2006 Jul 15;174(2):120-6. doi:10.1164/rccm.200601-084OC. Epub 2006 Apr 20. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16627867>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2662904/>)
- Kennedy MP, Omran H, Leigh MW, Dell S, Morgan L, Molina PL, Robinson BV, Minnix SL, Olbrich H, Severin T, Ahrens P, Lange L, Morillas HN, Noone PG, Zariwala MA, Knowles MR. Congenital heart disease and other heterotaxic defects in a large cohort of patients with primary ciliary dyskinesia. *Circulation*. 2007 Jun 5;115(22):2814-21. doi: 10.1161/CIRCULATIONAHA.106.649038. Epub 2007 May 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17515466>)
- Leigh MW, Pittman JE, Carson JL, Ferkol TW, Dell SD, Davis SD, Knowles MR, Zariwala MA. Clinical and genetic aspects of primary ciliary dyskinesia/Kartagener syndrome. *Genet Med*. 2009 Jul;11(7):473-87. doi:10.1097/GIM.0b013e3181a53562. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19606528>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3739704/>)
- Morillas HN, Zariwala M, Knowles MR. Genetic causes of bronchiectasis: primary ciliary dyskinesia. *Respiration*. 2007;74(3):252-63. doi: 10.1159/000101783. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17534128>)
- Shapiro AJ, Davis SD, Ferkol T, Dell SD, Rosenfeld M, Olivier KN, Sagel SD, Milla C, Zariwala MA, Wolf W, Carson JL, Hazucha MJ, Burns K, Robinson B, Knowles MR,

Leigh MW; Genetic Disorders of Mucociliary Clearance Consortium. Laterality defects other than situs inversus totalis in primary ciliary dyskinesia: insights into situs ambiguus and heterotaxy. *Chest*. 2014 Nov;146(5):1176-1186. doi:10.1378/chest.13-1704. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24577564>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4219335/>)

- Sutherland MJ, Ware SM. Disorders of left-right asymmetry: heterotaxy and situs inversus. *Am J Med Genet C Semin Med Genet*. 2009 Nov 15;151C(4):307-17. doi: 10.1002/ajmg.c.30228. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19876930>)
- Zariwala MA, Knowles MR, Leigh MW. Primary Ciliary Dyskinesia. 2007 Jan 24[updated 2019 Dec 5]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews*(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1122/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301301>)

**Last updated April 1, 2014**