

CARD11 gene

caspace recruitment domain family member 11

Normal Function

The *CARD11* gene provides instructions for making a protein involved in the function of immune system cells called lymphocytes, particularly certain types called T cells and B cells. These cells identify foreign substances such as bacteria, viruses, and fungi and defend the body against infection. When T or B cells recognize a foreign substance, the CARD11 protein is turned on (activated) and attaches (binds) to two other proteins, BCL10 and MALT1, to form the CBM signalosome complex. This complex in turn activates other protein complexes called nuclear factor-kappa-B (NF-κB) and mTOR complex 1 (mTORC1), which are important for cellular signaling. NF-κB and mTORC1 signaling direct the development and function of T and B cells so they can support an immune response against foreign invaders.

Health Conditions Related to Genetic Changes

Atopic dermatitis

At least five *CARD11* gene mutations have been identified in people with a skin disorder called atopic dermatitis (also known as atopic eczema). This condition is characterized by dry, itchy skin and red rashes. The word "atopic" indicates an association with allergies. While atopic dermatitis is not always due to an allergic reaction, it is commonly associated with other allergic disorders. People with atopic dermatitis caused by *CARD11* gene mutations often have additional allergic disorders, such as asthma and environmental (such as pollen) or food allergies. Many of these individuals also have recurrent infections due to problems with the immune system (immunodeficiency).

Atopic dermatitis is generally thought of as a complex condition that is influenced by multiple genetic and environmental factors, which each contribute only a small amount to the overall risk of developing the condition. However, *CARD11* gene mutations appear to cause atopic dermatitis without other factors. These mutations likely account for only a small percentage of cases of the condition.

A mutation in one of the two copies of the *CARD11* gene in each cell is sufficient to cause atopic dermatitis. These mutations result in the production of an altered protein that does not function normally. The altered protein produced from the mutated copy of the gene interferes with the normal protein produced from the non-mutated copy of the

gene (such mutations are described as "dominant-negative"), so the amount of functioning CARD11 protein in cells is reduced. These genetic changes are thought to prevent formation of the CBM signalosome complex, impairing signaling by NF-κB and mTORC1. Without these signals, T cells do not develop or function properly. The number of these cells is normal, but their response to foreign invaders is diminished, leading to recurrent infections.

It is not clear how the immune dysfunction caused by *CARD11* gene mutations leads to atopic dermatitis and allergic disorders. Atopic dermatitis is not initially caused by an allergic reaction, although sometimes substances that can cause allergic reactions (allergens) are thought to contribute to flare-ups of the rashes.

Omenn syndrome

MedlinePlus Genetics provides information about Omenn syndrome

Other disorders

CARD11 gene mutations can cause other immune system disorders. At least 3 mutations in the gene have been found to cause a type of severe combined immunodeficiency (SCID) known as immunodeficiency 11. SCID is a group of disorders characterized by an almost total lack of immune protection from foreign invaders. Immunodeficiency 11 is characterized by recurrent severe infections in the respiratory tract, particularly pneumonia caused by a fungus known as *Pneumocystis jirovecii*. Immunodeficiency 11 occurs when both copies of the *CARD11* gene are mutated. The mutations are described as "loss of function" because they lead to production of an abnormally short, nonfunctional CARD11 protein. The lack of CARD11 function diminishes T cells' ability to fight foreign invaders, despite normal numbers of the cells, leading to recurrent infections. Individuals with immunodeficiency 11 do not appear to have an increased risk of atopic dermatitis (described above) or allergies.

At least four mutations in the *CARD11* gene cause another immune cell disorder called B-cell expansion with NF-κB and T-cell anergy (BENTA). This condition is characterized by an excess of immune system cells called B cells (B-cell lymphocytosis), an increased risk of B-cell lymphoma, and susceptibility to infection. The mutations that cause BENTA are inherited and occur in one copy of the *CARD11* gene. Like those involved in B-cell and T-cell cancers (described above), the genetic changes that cause BENTA are "gain-of-function" mutations; they lead to production of an altered CARD11 protein that is always turned on, resulting in constant NF-κB signaling. Overactive NF-κB promotes the proliferation of B cells, which can lead to B-cell lymphoma. These abnormal cells, however, cannot respond to infections. T cells are also abnormal and unable to fight foreign invaders (a phenomenon known as anergy). These immune cell problems lead to the increased risk of infection in people with BENTA.

Cancers

Mutations in the *CARD11* gene are also associated with cancers of B cells (primarily diffuse large B-cell lymphoma) and T cells (adult T-cell leukemia/lymphoma). These

mutations are not inherited; instead, they arise during a person's lifetime and are found only in B or T cells that give rise to cancer. These genetic changes are called "gain-of-function" mutations because they lead to production of an altered CARD11 protein that is always turned on, even without recognition of foreign substances by the B or T cell. As a result, NF- κ B is constantly activated. Unregulated NF- κ B signaling allows these cells to grow and divide without control, contributing to the development of cancer.

Other Names for This Gene

- bcl10-interacting maguk protein 3
- BENTA
- BIMP3
- CARD-containing MAGUK protein 1
- carma 1
- CARMA1
- caspase recruitment domain-containing protein 11
- IMD11
- IMD11A
- PPBL

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of CARD11 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=84433\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=84433[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28CARD11%5BTIAB%5D%29+OR+%28caspase+recruitment+domain+family+member+11%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- OMENN SYNDROME (<https://omim.org/entry/603554>)
- CASPASE RECRUITMENT DOMAIN-CONTAINING PROTEIN 11; CARD11 (<https://omim.org/entry/607210>)
- B-CELL EXPANSION WITH NFKB AND T-CELL ANERGY; BENTA (<https://omim.org/entry/616452>)
- IMMUNODEFICIENCY 11A; IMD11A (<https://omim.org/entry/615206>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/84433>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=CARD11\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=CARD11[gene]))

References

- Dadi H, Jones TA, Merico D, Sharfe N, Ovadia A, Schejter Y, Reid B, Sun M, Vong L, Atkinson A, Lavi S, Pomerantz JL, Roifman CM. Combined immunodeficiency and atopy caused by a dominant negative mutation in caspase activation and recruitment domain family member 11 (CARD11). *J Allergy Clin Immunol*. 2018 May;141(5):1818-1830.e2. doi: 10.1016/j.jaci.2017.06.047. Epub 2017 Aug 19. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/28826773>)
- Greil J, Rausch T, Giese T, Bandapalli OR, Daniel V, Bekeredjian-Ding I, Stutz AM, Drees C, Roth S, Ruland J, Korbel JO, Kulozik AE. Whole-exome sequencing links caspase recruitment domain 11 (CARD11) inactivation to severe combined immunodeficiency. *J Allergy Clin Immunol*. 2013 May;131(5):1376-83.e3. doi:10.1016/j.jaci.2013.02.012. Epub 2013 Apr 3. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23561803>)
- Kogure Y, Kataoka K. Genetic alterations in adult T-cell leukemia/lymphoma. *Cancer Sci*. 2017 Sep;108(9):1719-1725. doi: 10.1111/cas.13303. Epub 2017 Jul 29. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/28627735>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5581529/>)
- Lamason RL, McCully RR, Lew SM, Pomerantz JL. Oncogenic CARD11 mutations induce hyperactive signaling by disrupting autoinhibition by the PKC-responsive inhibitory domain. *Biochemistry*. 2010 Sep 28;49(38):8240-50. doi:10.1021/bi101052d. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20799731>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2943563/>)
- Lenz G, Davis RE, Ngo VN, Lam L, George TC, Wright GW, Dave SS, Zhao H, Xu W, Rosenwald A, Ott G, Muller-Hermelink HK, Gascoyne RD, Connors JM, Rimsza LM, Campo E, Jaffe ES, Delabie J, Smeland EB, Fisher RI, Chan WC, Staudt LM. Oncogenic CARD11 mutations in human diffuse large B cell lymphoma. *Science*. 2008 Mar 21;319(5870):1676-9. doi: 10.1126/science.1153629. Epub 2008 Mar 6. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18323416>)
- Ma CA, Stinson JR, Zhang Y, Abbott JK, Weinreich MA, Hauk PJ, Reynolds PR, Lyons JJ, Nelson CG, Ruffo E, Dorjbal B, Glauzy S, Yamakawa N, Arjunaraja S, Voss K, Stoddard J, Niemela J, Zhang Y, Rosenzweig SD, McElwee JJ, DiMaggio T, Matthews HF, Jones N, Stone KD, Palma A, Oleastro M, Prieto E, Bernasconi AR, Dubra G, Danielian S, Zaiat J, Marti MA, Kim B, Cooper MA, Romberg N, Meffre E, Gelfand EW, Snow AL, Milner JD. Germline hypomorphic CARD11 mutations in severe atopic disease. *Nat Genet*. 2017 Aug;49(8):1192-1201. doi: 10.1038/ng.3898. Epub 2017 Jun 19. Erratum In: *Nat Genet*. 2017 Oct 27;49(11):1661. Citation on

PubMed (<https://pubmed.ncbi.nlm.nih.gov/28628108>)

- Shi JH, Sun SC. TCR signaling to NF-kappaB and mTORC1: Expanding roles of theCARMA1 complex. *Mol Immunol*. 2015 Dec;68(2 Pt C):546-57. doi:10.1016/j.molimm.2015.07.024. Epub 2015 Aug 8. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/26260210>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4679546/>)
- Snow AL, Xiao W, Stinson JR, Lu W, Chaigne-Delalande B, Zheng L, Pittaluga S, Matthews HF, Schmitz R, Jhavar S, Kuchen S, Kardava L, Wang W, Lamborn IT, JingH, Raffeld M, Moir S, Fleisher TA, Staudt LM, Su HC, Lenardo MJ. Congenital Bcell lymphocytosis explained by novel germline CARD11 mutations. *J Exp Med*. 2012Nov 19;209(12):2247-61. doi: 10.1084/jem.20120831. Epub 2012 Nov 5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23129749>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3501355/>)
- Thome M, Charton JE, Pelzer C, Hailfinger S. Antigen receptor signaling toNF-kappaB via CARMA1, BCL10, and MALT1. *Cold Spring Harb Perspect Biol*. 2010Sep;2(9):a003004. doi: 10.1101/cshperspect.a003004. Epub 2010 Aug 4. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20685844>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2926749/>)
- Turvey SE, Durandy A, Fischer A, Fung SY, Geha RS, Gewies A, Giese T, Greil J, Keller B, McKinnon ML, Neven B, Rozmus J, Ruland J, Snow AL, Stepensky P, WarnatzK. The CARD11-BCL10-MALT1 (CBM) signalosome complex: Stepping into the limelightof human primary immunodeficiency. *J Allergy Clin Immunol*. 2014Aug; 134(2):276-84. doi: 10.1016/j.jaci.2014.06.015. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25087226>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4167767/>)

Genomic Location

The *CARD11* gene is found on chromosome 7 (<https://medlineplus.gov/genetics/chromosome/7/>).

Last updated October 1, 2017