

COL17A1 gene

collagen type XVII alpha 1 chain

Normal Function

The *COL17A1* gene provides instructions for making a protein that is used to assemble type XVII collagen. Collagens are a family of proteins that strengthen and support connective tissues, such as skin, bone, tendons, and ligaments, throughout the body. In particular, type XVII collagen plays an essential role in strengthening and stabilizing the skin.

The protein produced from the *COL17A1* gene is known as a pro- α 1(XVII) chain. Three identical pro- α 1(XVII) chains twist together to form a triple-stranded, ropelike molecule known as a procollagen. Procollagen molecules are released from the cell and processed by enzymes to remove extra protein segments from the ends. Once these molecules are processed, they arrange themselves into long, thin bundles of mature type XVII collagen.

Type XVII collagen is a major component of hemidesmosomes, which are microscopic structures on the inner surface of the top layer of skin (the epidermis). These structures help to anchor the epidermis to underlying layers of skin. Type XVII collagen is critical for the stability of hemidesmosomes, and therefore it plays an important role in holding the layers of skin together.

Health Conditions Related to Genetic Changes

Junctional epidermolysis bullosa

More than 100 mutations in the *COL17A1* gene have been identified in people with junctional epidermolysis bullosa (JEB). Most of these mutations add or remove several DNA building blocks (nucleotides) in the *COL17A1* gene or create a premature stop signal in the instructions for making the pro- α 1(XVII) chain. These changes reduce the amount of functional type XVII collagen in the skin. Without enough of this collagen, the epidermis is only weakly attached to underlying layers of skin. Friction or other minor trauma (such as rubbing or scratching) can cause the skin layers to separate, leading to the formation of blisters.

Most *COL17A1* gene mutations cause the milder form of junctional epidermolysis bullosa, known as JEB generalized intermediate. Affected individuals experience

blistering, but it may be limited to the hands, feet, knees, and elbows and often improves after the newborn period. A few individuals with mutations in the *COL17A1* gene have had the more serious form of the disorder, JEB generalized severe. Infants with JEB generalized severe develop widespread blistering that causes life-threatening complications.

Other Names for This Gene

- alpha 1 type XVII collagen
- BA16H23.2
- BP180
- BPAG2
- bullous pemphigoid antigen 2 (180kD)
- COHA1_HUMAN
- collagen type XVII alpha 1
- collagen XVII, alpha-1 polypeptide
- collagen, type XVII, alpha 1
- FLJ60881
- KIAA0204
- LAD-1

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of COL17A1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=1308\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=1308[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28COL17A1%5BTIAB%5D%29+OR+%28%28type+XVII+collagen+%5Btiab%5D+AND+alpha+1+%5Btiab%5D%29+OR+%28BP180%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- COLLAGEN, TYPE XVII, ALPHA-1; COL17A1 (<https://omim.org/entry/113811>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/1308>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=COL17A1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=COL17A1[gene]))

References

- Bauer JW, Lanschuetzer C. Type XVII collagen gene mutations in junctionalepidermolysis bullosa and prospects for gene therapy. *Clin Exp Dermatol*. 2003Jan;28(1):53-60. doi: 10.1046/j.1365-2230.2003.01192.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12558632>)
- Franzke CW, Has C, Schulte C, Huilaja L, Tasanen K, Aumailley M, Bruckner-Tuderman L. C-terminal truncation impairs glycosylation of transmembranecollagen XVII and leads to intracellular accumulation. *J Biol Chem*. 2006 Oct6;281(40):30260-8. doi: 10.1074/jbc.M604464200. Epub 2006 Aug 9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16899459>)
- Fu CL, Giudice GJ, Van den Bergh F. Protein structural analysis of BP180mutant isoforms linked to non-Herlitz junctional epidermolysis bullosa. *J InvestDermatol*. 2006 Jan;126(1):232-4. doi: 10.1038/sj.jid.5700024. No abstractavailable. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16417243>)
- Gatalica B, Pulkkinen L, Li K, Kuokkanen K, Ryyanen M, McGrath JA, Uitto J. Cloning of the human type XVII collagen gene (COL17A1), and detection of novelmutations in generalized atrophic benign epidermolysis bullosa. *Am J Hum Genet*. 1997 Feb;60(2):352-65. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9012408>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1712405/>)
- Nakamura H, Sawamura D, Goto M, Nakamura H, Kida M, Ariga T, Sakiyama Y, Tomizawa K, Mitsui H, Tamaki K, Shimizu H. Analysis of the COL17A1 in non-Herlitzjunctional epidermolysis bullosa and amelogenesis imperfecta. *Int J Mol Med*. 2006Aug;18(2):333-7. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16820943>)
- Pasmooij AM, Pas HH, Jansen GH, Lemmink HH, Jonkman MF. Localized andgeneralized forms of blistering in junctional epidermolysis bullosa due toCOL17A1 mutations in the Netherlands. *Br J Dermatol*. 2007 May;156(5):861-70. doi:10.1111/j.1365-2133.2006.07730.x. Epub 2007 Jan 30. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17263807>)
- Pulkkinen L, Uitto J. Mutation analysis and molecular genetics ofepidermolysis bullosa. *Matrix Biol*. 1999 Feb;18(1):29-42. doi:10.1016/s0945-053x(98)00005-5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10367729>)
- Ruzzi L, Pas H, Posteraro P, Mazzanti C, Didona B, Owaribe K, Meneguzzi G, Zambruno G, Castiglia D, Alessio M. A homozygous nonsense mutation in type XVIIcollagen gene (COL17A1) uncovers an alternatively spliced mRNA accounting for anunusually mild form of non-Herlitz junctional epidermolysis bullosa. *J InvestDermatol*. 2001 Jan;116(1):182-7. doi: 10.1046/j.1523-1747.2001.00229.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11168815>)
- Varki R, Sadowski S, Pfendner E, Uitto J. Epidermolysis bullosa. I.

Molecular genetics of the junctional and hemidesmosomal variants. J Med Genet. 2006 Aug;43(8):52. doi: 10.1136/jmg.2005.039685. Epub 2006 Feb 10. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16473856>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2564586/>)

Genomic Location

The *COL17A1* gene is found on chromosome 10 (<https://medlineplus.gov/genetics/chromosome/10/>).

Last updated July 1, 2019