

DNAJC5 gene

DnaJ heat shock protein family (Hsp40) member C5

Normal Function

The *DNAJC5* gene provides instructions for making a protein called cysteine string protein alpha (CSP α). This protein is found near nerve cells in the brain, where it plays a role in the transmission of nerve impulses. Specifically, CSP α is part of a group (complex) of proteins that is found on the membrane of sac-like structures called synaptic vesicles. Synaptic vesicles are found close to the ends of nerve cells and contain chemical messengers that transmit signals from one nerve cell to another. CSP α is involved in recycling proteins that are involved in nerve impulse transmission by re-folding misshapen proteins so that they can be used in additional transmissions.

Health Conditions Related to Genetic Changes

CLN4 disease

At least two mutations in the *DNAJC5* gene have been found to cause CLN4 disease. CLN4 disease is an inherited disorder that primarily affects the nervous system. This condition usually begins in adulthood with problems with movement and intellectual function that worsen over time.

One of the *DNAJC5* gene mutations replaces the protein building block (amino acid) leucine with the amino acid arginine at position 115 in the CSP α protein (written as L115R). The other mutation deletes the amino acid leucine at position 116 in the protein (written as L116del). Affected individuals have one of these mutations in one copy of the *DNAJC5* gene in each cell, which leads to the production of an altered protein that cannot associate with the membrane of synaptic vesicles. The resulting reduction in protein recycling leads to a shortage (deficiency) of functional proteins, which impairs the efficiency of nerve impulse transmission. The abnormal CSP α protein may also bind to the functional CSP α protein that is produced from the normal copy of the *DNAJC5* gene and prevent it from associating with synaptic vesicles, further impairing impulse transmission. Without communication between nerve cells, neurological functions are impaired, contributing to the features of CLN4 disease.

CLN4 disease is characterized by the accumulation of proteins and other substances in lysosomes, which are compartments in the cell that digest and recycle materials. These accumulations occur in cells throughout the body; however, nerve cells seem to be

particularly vulnerable to their effects. The accumulations can cause cell damage leading to cell death. The progressive death of nerve cells in the brain and other tissues contributes to the decline of neurological function in CLN4 disease. However, it is unclear how mutations in the *DNAJC5* gene are involved in the buildup of substances in lysosomes.

Other Names for This Gene

- CLN4
- CLN4B
- CSP
- cysteine string protein alpha
- DnaJ (Hsp40) homolog, subfamily C, member 5
- dnaJ homolog subfamily C member 5
- DNAJC5A
- DNJC5_HUMAN
- FLJ00118
- FLJ13070

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of DNAJC5 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=80331\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=80331[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28DNAJC5%5BTIAB%5D%29+OR+%28CLN4%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D%29%29%29>)

Catalog of Genes and Diseases from OMIM

- DNAJ/HSP40 HOMOLOG, SUBFAMILY C, MEMBER 5; DNAJC5 (<https://omim.org/entry/611203>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/80331>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=DNAJC5\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=DNAJC5[gene]))

References

- Benitez BA, Alvarado D, Cai Y, Mayo K, Chakraverty S, Norton J, Morris JC, Sands MS, Goate A, Cruchaga C. Exome-sequencing confirms DNAJC5 mutations as cause of adult neuronal ceroid-lipofuscinosis. PLoS One. 2011;6(11):e26741. doi: 10.1371/journal.pone.0026741. Epub 2011 Nov 4. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22073189>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3208569/>)
- Cadieux-Dion M, Andermann E, Lachance-Touchette P, Ansorge O, Meloche C, Barnabe A, Kuzniecky RI, Andermann F, Faught E, Leonberg S, Damiano JA, Berkovic SF, Rouleau GA, Cossette P. Recurrent mutations in DNAJC5 cause autosomal dominant Kufs disease. Clin Genet. 2013 Jun;83(6):571-5. doi: 10.1111/cge.12020. Epub 2012 Nov 7. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22978711>)
- Carcel-Trullols J, Kovacs AD, Pearce DA. Cell biology of the NCL proteins: What they do and don't do. Biochim Biophys Acta. 2015 Oct;1852(10 Pt B):2242-55. doi: 10.1016/j.bbadis.2015.04.027. Epub 2015 May 8. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25962910>)
- Henderson MX, Wirak GS, Zhang YQ, Dai F, Ginsberg SD, Dolzhanskaya N, Staropoli JF, Nijssen PC, Lam TT, Roth AF, Davis NG, Dawson G, Velinov M, Chandra SS. Neuronal ceroid lipofuscinosis with DNAJC5/CSPalpha mutation has PPT1 pathology and exhibit aberrant protein palmitoylation. Acta Neuropathol. 2016 Apr;131(4):621-37. doi: 10.1007/s00401-015-1512-2. Epub 2015 Dec 10. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/26659577>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4791186/>)
- Tobaben S, Thakur P, Fernandez-Chacon R, Sudhof TC, Rettig J, Stahl B. A trimeric protein complex functions as a synaptic chaperone machine. Neuron. 2001 Sep 27;31(6):987-99. doi: 10.1016/s0896-6273(01)00427-5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11580898>)
- Velinov M, Dolzhanskaya N, Gonzalez M, Powell E, Konidari I, Hulme W, Staropoli JF, Xin W, Wen GY, Barone R, Coppel SH, Sims K, Brown WT, Zuchner S. Mutations in the gene DNAJC5 cause autosomal dominant Kufs disease in a proportion of cases: study of the Parry family and 8 other families. PLoS One. 2012;7(1):e29729. doi: 10.1371/journal.pone.0029729. Epub 2012 Jan 3. Erratum In: PLoS One. 2012;7(9). doi:10.1371/annotation/26d7eb64-ccd2-41db-b1aa-7cdc8c1eff95. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22235333>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3250487/>)

Genomic Location

The *DNAJC5* gene is found on chromosome 20 (<https://medlineplus.gov/genetics/chromosome/20/>).

Last updated January 1, 2017