

EDA gene

ectodysplasin A

Normal Function

The *EDA* gene provides instructions for making a protein called ectodysplasin A. This protein is part of a signaling pathway that plays an important role in development before birth. Specifically, it is critical for interactions between two embryonic cell layers called the ectoderm and the mesoderm. In the early embryo, these cell layers form the basis for many of the body's organs and tissues. Ectoderm-mesoderm interactions are essential for the formation of several structures that arise from the ectoderm, including the skin, hair, nails, teeth, and sweat glands.

One version of the ectodysplasin A protein, known as ectodysplasin A1, interacts with a protein called the ectodysplasin A receptor (produced from the *EDAR* gene). On the cell surface, ectodysplasin A1 attaches to this receptor like a key in a lock. When these two proteins are connected, they trigger a series of chemical signals that affect cell activities such as division, growth, and maturation. Starting before birth, this signaling pathway controls the formation of ectodermal structures such as hair follicles, sweat glands, and teeth.

Health Conditions Related to Genetic Changes

Hypohidrotic ectodermal dysplasia

More than 300 mutations in the *EDA* gene have been found to cause hypohidrotic ectodermal dysplasia, the most common form of ectodermal dysplasia. Starting before birth, ectodermal dysplasias result in the abnormal development of the skin, hair, nails, teeth, and sweat glands. Hypohidrotic ectodermal dysplasia is characterized by a reduced ability to sweat (hypohidrosis), sparse scalp and body hair (hypotrichosis), and several missing teeth (hypodontia) or teeth that are malformed. *EDA* gene mutations are the most frequent cause of hypohidrotic ectodermal dysplasia, accounting for more than half of all cases.

Some mutations in the *EDA* gene change single DNA building blocks (base pairs), whereas other mutations insert or delete a larger section of DNA. These changes lead to the production of a nonfunctional version of the ectodysplasin A1 protein. A shortage of functional ectodysplasin A1 prevents the protein from interacting effectively with its receptor, which impairs chemical signaling needed for interactions between the

ectoderm and the mesoderm in early development. Without these signals, hair follicles, teeth, sweat glands, and other ectodermal structures do not form properly, which leads to the characteristic features of hypohidrotic ectodermal dysplasia.

Other disorders

EDA gene mutations have also been reported in some people with a condition called nonsyndromic tooth agenesis. This condition causes one or more teeth not to form. Although missing teeth is a common feature of ectodermal dysplasias, "nonsyndromic" suggests that in these cases tooth agenesis occurs without the other signs and symptoms of those conditions. It is unclear why the effects of some mutations in this gene appear to be limited to tooth development, while other mutations affect the formation of multiple ectodermal tissues.

Other Names for This Gene

- Ectodermal dysplasia protein
- ectodysplasin
- ectodysplasin-A
- ED1
- ED1-A1
- EDA-A1
- EDA-A2
- EDA1
- EDA_HUMAN
- HED
- XHED
- XLHED

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of EDA ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=1896\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=1896[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28ectodysplasin+A%5BTIAB%5D%29+OR+%28%28ectodysplasin-A%5BTIAB%5D%29+OR+%28EDA1%5BTIAB%5D%29%29+OR+%28%28EDA%5BTI%5D%29+AND+%28ectodermal+dysplasia%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D>)

Catalog of Genes and Diseases from OMIM

- ECTODYSPLASIN A; EDA (<https://omim.org/entry/300451>)
- TOOTH AGENESIS, SELECTIVE, X-LINKED, 1; STHAGX1 (<https://omim.org/entry/313500>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/1896>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=EDA\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=EDA[gene]))

References

- Cluzeau C, Hadj-Rabia S, Jambou M, Mansour S, Guigue P, Masmoudi S, Bal E, Chassaing N, Vincent MC, Viot G, Clauss F, Maniere MC, Toupenay S, Le Merrer M, Lyonnet S, Cormier-Daire V, Amiel J, Faivre L, de Prost Y, Munnich A, BonnefontJP, Bodemer C, Smahi A. Only four genes (EDA1, EDAR, EDARADD, and WNT10A) accountfor 90% of hypohidrotic/anhidrotic ectodermal dysplasia cases. Hum Mutat. 2011Jan;32(1):70-2. doi: 10.1002/humu.21384. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20979233>)
- Han D, Gong Y, Wu H, Zhang X, Yan M, Wang X, Qu H, Feng H, Song S. Novel EDAmutation resulting in X-linked non-syndromic hypodontia and the pattern ofEDA-associated isolated tooth agenesis. Eur J Med Genet. 2008Nov-Dec;51(6):536-46. doi: 10.1016/j.ejmg.2008.06.002. Epub 2008 Jul 9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18657636>)
- Kowalczyk-Quintas C, Schneider P. Ectodysplasin A (EDA) - EDA receptorgnalling and its pharmacological modulation. Cytokine Growth Factor Rev. 2014Apr;25(2):195-203. doi: 10.1016/j.cytogfr.2014.01.004. Epub 2014 Jan 23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24508088>)
- Song S, Han D, Qu H, Gong Y, Wu H, Zhang X, Zhong N, Feng H. EDA genemutations underlie non-syndromic oligodontia. J Dent Res. 2009 Feb;88(2):126-31.doi: 10.1177/0022034508328627. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19278982>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3317984/>)
- Tarpey P, Pemberton TJ, Stockton DW, Das P, Ninis V, Edkins S, Andrew FutrealP, Wooster R, Kamath S, Nayak R, Stratton MR, Patel PI. A novel Gln358Glu mutation in ectodysplasin A associated with X-linked dominant incisor hypodontia.Am J Med Genet A. 2007 Feb 15;143(4):390-4. doi: 10.1002/ajmg.a.31567. Noabstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17256800>)
- Vincent MC, Biancalana V, Ginisty D, Mandel JL, Calvas P. Mutational spectrumof the ED1 gene in X-linked hypohidrotic ectodermal dysplasia. Eur J Hum Genet.2001 May;9(5):355-63. doi: 10.1038/sj.ejhg.5200635. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11378824>)
- Wisniewski SA, Kobiela A, Trzeciak WH, Kobiela K. Recent advances

in understanding of the molecular basis of anhidrotic ectodermal dysplasia: discovery of a ligand, ectodysplasin A and its two receptors. J Appl Genet. 2002;43(1):97-107. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12084975>)

- Wohlfart S, Hammersen J, Schneider H. Mutational spectrum in 101 patients with hypohidrotic ectodermal dysplasia and breakpoint mapping in independent cases of rare genomic rearrangements. J Hum Genet. 2016 Oct;61(10):891-897. doi: 10.1038/jhg.2016.75. Epub 2016 Jun 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27305980>)
- Wright JT, Grange DK, Fete M. Hypohidrotic Ectodermal Dysplasia. 2003 Apr 28 [updated 2022 Oct 27]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. GeneReviews(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1112/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301291>)

Genomic Location

The *EDA* gene is found on the X chromosome (<https://medlineplus.gov/genetics/chromosome/x/>).

Last updated November 1, 2018