

## EDAR gene

ectodysplasin A receptor

### Normal Function

The *EDAR* gene provides instructions for making a protein called the ectodysplasin A receptor. This protein is part of a signaling pathway that plays an important role in development before birth. Specifically, it is critical for interactions between two embryonic cell layers called the ectoderm and the mesoderm. In the early embryo, these cell layers form the basis for many of the body's organs and tissues. Ectoderm-mesoderm interactions are essential for the formation of several structures that arise from the ectoderm, including the skin, hair, nails, teeth, and sweat glands.

The ectodysplasin A receptor interacts with a protein called ectodysplasin A1 (produced from the *EDA* gene). On the cell surface, ectodysplasin A1 attaches to this receptor like a key in a lock. When these two proteins are connected, they trigger a series of chemical signals that affect cell activities such as division, growth, and maturation. Starting before birth, this signaling pathway controls the formation of ectodermal structures such as hair follicles, sweat glands, and teeth.

Studies suggest that common variations (polymorphisms) in the *EDAR* gene are associated with the thickness and straightness of scalp hair, particularly in East Asian populations. *EDAR* appears to be one of many genes that influence these hair traits.

### Health Conditions Related to Genetic Changes

#### Hypohidrotic ectodermal dysplasia

More than 50 mutations in the *EDAR* gene have been found to cause hypohidrotic ectodermal dysplasia, the most common form of ectodermal dysplasia. Starting before birth, ectodermal dysplasias result in the abnormal development of the skin, hair, nails, teeth, and sweat glands. Hypohidrotic ectodermal dysplasia is characterized by a reduced ability to sweat (hypohidrosis), sparse scalp and body hair (hypotrichosis), and several missing teeth (hypodontia) or teeth that are malformed. *EDAR* gene mutations account for about 10 percent of all cases of hypohidrotic ectodermal dysplasia.

Most of the *EDAR* gene mutations associated with hypohidrotic ectodermal dysplasia change a single protein building block (amino acid) in the receptor protein. Some of the mutations that cause this condition lead to the production of an abnormal version of the

ectodysplasin A receptor. Other mutations prevent cells from producing any functional receptor. All of these genetic changes prevent the receptor from interacting with ectodysplasin A1, which impairs chemical signaling needed for interactions between the ectoderm and the mesoderm in early development. Without these signals, hair follicles, teeth, sweat glands, and other ectodermal structures do not form properly, which leads to the characteristic features of hypohidrotic ectodermal dysplasia.

### Other disorders

*EDAR* gene mutations have also been reported in some people with a condition called nonsyndromic tooth agenesis. This condition causes one or more teeth not to form. Although missing teeth is a common feature of ectodermal dysplasias, "nonsyndromic" suggests that in these cases tooth agenesis occurs without the other signs and symptoms of those conditions. It is unclear why the effects of some mutations in this gene appear to be limited to tooth development, while other mutations affect the formation of multiple ectodermal tissues.

### **Other Names for This Gene**

- DL
- ectodysplasin 1, anhidrotic receptor
- ectodysplasin A1 isoform receptor
- ectodysplasin receptor
- ED1R
- ED3
- ED5
- EDA-A1R
- EDA1R
- EDA3
- EDAR\_HUMAN

### **Additional Information & Resources**

#### Tests Listed in the Genetic Testing Registry

- Tests of EDAR ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=10913\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=10913[geneid]))

#### Scientific Articles on PubMed

- PubMed ([https://pubmed.ncbi.nlm.nih.gov/?term=%28%28ectodysplasin+A+receptor%5BTIAB%5D%29+OR+%28ectodysplasin+A1+receptor%5BTIAB%5D%29%29+OR+%28%28EDAR%5BTIAB%5D%29+NOT+%28Alzheimer\\*%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D](https://pubmed.ncbi.nlm.nih.gov/?term=%28%28ectodysplasin+A+receptor%5BTIAB%5D%29+OR+%28ectodysplasin+A1+receptor%5BTIAB%5D%29%29+OR+%28%28EDAR%5BTIAB%5D%29+NOT+%28Alzheimer*%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D))

### Catalog of Genes and Diseases from OMIM

- ECTODYSPLASIN A RECEPTOR; EDAR (<https://omim.org/entry/604095>)
- HAIR MORPHOLOGY 1; HRM1 (<https://omim.org/entry/612630>)

### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/10913>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=EDAR\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=EDAR[gene]))

### **References**

- Arte S, Parmanen S, Pirinen S, Alaluusua S, Nieminen P. Candidate gene analysis of tooth agenesis identifies novel mutations in six genes and suggests significant role for WNT and EDA signaling and allele combinations. PLoS One. 2013 Aug 22;8(8):e73705. doi: 10.1371/journal.pone.0073705. eCollection 2013. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23991204>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3750013/>)
- Azeem Z, Naqvi SK, Ansar M, Wali A, Naveed AK, Ali G, Hassan MJ, Tariq M, Basit S, Ahmad W. Recurrent mutations in functionally-related EDA and EDAR genes underlie X-linked isolated hypodontia and autosomal recessive hypohidrotic ectodermal dysplasia. Arch Dermatol Res. 2009 Sep;301(8):625-9. doi: 10.1007/s00403-009-0975-1. Epub 2009 Jun 24. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19551394>)
- Bashyam MD, Chaudhary AK, Reddy EC, Reddy V, Acharya V, Nagarajaram HA, Devi AR, Bashyam L, Dalal AB, Gupta N, Kabra M, Agarwal M, Phadke SR, Tainwala R, Kumar R, Hariharan SV. A founder ectodysplasin A receptor (EDAR) mutation results in a high frequency of the autosomal recessive form of hypohidrotic ectodermal dysplasia in India. Br J Dermatol. 2012 Apr;166(4):819-29. doi:10.1111/j.1365-2133.2011.10707.x. Epub 2012 Mar 5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22032522>)
- Cluzeau C, Hadj-Rabia S, Jambou M, Mansour S, Guigue P, Masmoudi S, Bal E, Chassaing N, Vincent MC, Viot G, Clauss F, Maniere MC, Toupenay S, Le Merrer M, Lyonnet S, Cormier-Daire V, Amiel J, Faivre L, de Prost Y, Munnich A, Bonnefont JP, Bodemer C, Smahi A. Only four genes (EDA1, EDAR, EDARADD, and WNT10A) account for 90% of hypohidrotic/anhydrotic ectodermal dysplasia cases. Hum Mutat. 2011 Jan;32(1):70-2. doi: 10.1002/humu.21384. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20979233>)
- Monreal AW, Ferguson BM, Headon DJ, Street SL, Overbeek PA, Zonana J. Mutations in the human homologue of mouse dl cause autosomal recessive and dominant hypohidrotic ectodermal dysplasia. Nat Genet. 1999 Aug;22(4):366-9. doi:10.1038/11937. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10431241>)

- Naeem M, Muhammad D, Ahmad W. Novel mutations in the EDAR gene in two Pakistani consanguineous families with autosomal recessive hypohidrotic ectodermal dysplasia. *Br J Dermatol.* 2005 Jul;153(1):46-50. doi:10.1111/j.1365-2133.2005.06642.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16029325>)
- Wisniewski SA, Kobiela A, Trzeciak WH, Kobiela K. Recent advances in understanding of the molecular basis of anhidrotic ectodermal dysplasia: discovery of a ligand, ectodysplasin A and its two receptors. *J Appl Genet.* 2002;43(1):97-107. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12084975>)
- Wohlfart S, Hammersen J, Schneider H. Mutational spectrum in 101 patients with hypohidrotic ectodermal dysplasia and breakpoint mapping in independent cases of rare genomic rearrangements. *J Hum Genet.* 2016 Oct;61(10):891-897. doi: 10.1038/jhg.2016.75. Epub 2016 Jun 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27305980>)
- Wright JT, Grange DK, Fete M. Hypohidrotic Ectodermal Dysplasia. 2003 Apr 28 [updated 2022 Oct 27]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews*(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1112/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301291>)
- Wu S, Tan J, Yang Y, Peng Q, Zhang M, Li J, Lu D, Liu Y, Lou H, Feng Q, Lu Y, Guan Y, Zhang Z, Jiao Y, Sabeti P, Krutmann J, Tang K, Jin L, Xu S, Wang S. Genome-wide scans reveal variants at EDAR predominantly affecting hair straightness in Han Chinese and Uyghur populations. *Hum Genet.* 2016 Nov;135(11):1279-1286. doi: 10.1007/s00439-016-1718-y. Epub 2016 Aug 3. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27487801>)
- Zeng B, Zhao Q, Li S, Lu H, Lu J, Ma L, Zhao W, Yu D. Novel EDA or EDAR Mutations Identified in Patients with X-Linked Hypohidrotic Ectodermal Dysplasia or Non-Syndromic Tooth Agenesis. *Genes (Basel).* 2017 Oct 5;8(10):259. doi:10.3390/genes8100259. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/28981473>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5664109/>)

## Genomic Location

The *EDAR* gene is found on chromosome 2 (<https://medlineplus.gov/genetics/chromosome/2/>).

**Last updated November 1, 2018**