

ENG gene

endoglin

Normal Function

The *ENG* gene provides instructions for making a protein called endoglin. This protein is found on the surface of cells, especially in the lining of developing arteries. It forms a complex with growth factors and other proteins involved in the development of blood vessels. In particular, this complex is involved in the specialization of new blood vessels into arteries or veins.

Health Conditions Related to Genetic Changes

Hereditary hemorrhagic telangiectasia

Dozens of mutations in the *ENG* gene have been found to cause hereditary hemorrhagic telangiectasia type 1. Many *ENG* gene mutations substitute one protein building block (amino acid) for another amino acid in the endoglin protein, which impairs the protein's function. Other mutations prevent production of the endoglin protein or result in an abnormally small protein that cannot function. The shortage of functional endoglin appears to interfere with the development of boundaries between arteries and veins, resulting in the signs and symptoms of hereditary hemorrhagic telangiectasia type 1.

Pulmonary arterial hypertension

MedlinePlus Genetics provides information about Pulmonary arterial hypertension

Other Names for This Gene

- CD105
- EGLN_HUMAN
- END
- endoglin (Osler-Rendu-Weber syndrome 1)
- endoglin precursor
- HHT1
- ORW

- ORW1
- Transforming Growth Factor P Receptor III

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of ENG ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2022\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2022[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28ENG%5BTIAB%5D%29+OR+%28endoglin%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29%29%29>)

Catalog of Genes and Diseases from OMIM

- ENDOGLIN; ENG (<https://omim.org/entry/131195>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/2022>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=ENG\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=ENG[gene]))

References

- Azuma H. Genetic and molecular pathogenesis of hereditary hemorrhagictelangiectasia. J Med Invest. 2000 Aug;47(3-4):81-90. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11019486>)
- Conley BA, Koleva R, Smith JD, Kacer D, Zhang D, Bernabeu C, Vary CP. Endoglincontrols cell migration and composition of focal adhesions: function of thecytosolic domain. J Biol Chem. 2004 Jun 25;279(26):27440-9. doi:10.1074/jbc.M312561200. Epub 2004 Apr 14. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15084601>)
- Karabegovic A, Shinawi M, Cymerman U, Letarte M. No live individual homozygousfor a novel endoglin mutation was found in a consanguineous Arab family withhereditary haemorrhagic telangiectasia. J Med Genet. 2004 Nov;41(11):e119. doi:10.1136/jmg.2004.022079. No abstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15520401>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735618/>)
- Lebrin F, Deckers M, Bertolino P, Ten Dijke P. TGF-beta receptor function inthe

endothelium. *Cardiovasc Res*. 2005 Feb 15;65(3):599-608. doi:10.1016/j.cardiores.2004.10.036. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15664386>)

- Lesca G, Plauchu H, Coulet F, Lefebvre S, Plessis G, Odent S, Riviere S, Leheup B, Goizet C, Carette MF, Cordier JF, Pinson S, Soubrier F, Calender A, Giraud S; French Rendu-Osler Network. Molecular screening of ALK1/ACVRL1 and ENG genes in hereditary hemorrhagic telangiectasia in France. *Hum Mutat*. 2004 Apr; 23(4):289-99. doi: 10.1002/humu.20017. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15024723>)
- Letteboer TG, Zewald RA, Kamping EJ, de Haas G, Mager JJ, Snijder RJ, Lindhout D, Hennekam FA, Westermann CJ, Ploos van Amstel JK. Hereditary hemorrhagic telangiectasia: ENG and ALK-1 mutations in Dutch patients. *Hum Genet*. 2005 Jan; 116(1-2):8-16. doi: 10.1007/s00439-004-1196-5. Epub 2004 Oct 23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15517393>)
- Pawlikowska L, Tran MN, Achrol AS, Ha C, Burchard E, Choudhry S, Zaroff J, Lawton MT, Castro R, McCulloch CE, Marchuk D, Kwok PY, Young WL; UCSF BAVM Study Project. Polymorphisms in transforming growth factor-beta-related genes ALK1 and ENG are associated with sporadic brain arteriovenous malformations. *Stroke*. 2005 Oct; 36(10):2278-80. doi: 10.1161/01.STR.0000182253.91167.f. Epub 2005 Sep 22. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16179574>)
- Sanz-Rodriguez F, Fernandez-L A, Zarrabeitia R, Perez-Molino A, Ramirez JR, Coto E, Bernabeu C, Botella LM. Mutation analysis in Spanish patients with hereditary hemorrhagic telangiectasia: deficient endoglin up-regulation in activated monocytes. *Clin Chem*. 2004 Nov; 50(11):2003-11. doi:10.1373/clinchem.2004.035287. Epub 2004 Sep 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15375013>)
- van den Driesche S, Mummery CL, Westermann CJ. Hereditary hemorrhagic telangiectasia: an update on transforming growth factor beta signaling in vasculogenesis and angiogenesis. *Cardiovasc Res*. 2003 Apr 1; 58(1):20-31. doi: 10.1016/s0008-6363(02)00852-0. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12667943>)

Genomic Location

The *ENG* gene is found on chromosome 9 (<https://medlineplus.gov/genetics/chromosome/9/>).

Last updated February 1, 2007