

EXT1 gene

exostosin glycosyltransferase 1

Normal Function

The *EXT1* gene provides instructions for producing a protein called exostosin-1. This protein is found in a cell structure called the Golgi apparatus, which modifies newly produced enzymes and other proteins. In the Golgi apparatus, exostosin-1 attaches (binds) to another protein, exostosin-2, to form a complex that modifies heparan sulfate. Heparan sulfate is a complex of sugar molecules (a polysaccharide) that is added to proteins to form proteoglycans, which are proteins attached to several sugars. Heparan sulfate is involved in regulating a variety of body processes including blood clotting and the formation of blood vessels (angiogenesis). It also has a role in the spreading (metastasis) of cancer cells.

Health Conditions Related to Genetic Changes

Hereditary multiple osteochondromas

About 480 mutations in the *EXT1* gene have been identified in people with hereditary multiple osteochondromas type 1, a condition in which people develop multiple benign (noncancerous) bone tumors called osteochondromas. Most of these mutations are known as "loss-of-function" mutations because they prevent any functional exostosin-1 protein from being made. The loss of functional exostosin-1 protein prevents it from forming a complex with the exostosin-2 protein and adding heparan sulfate to proteins. It is unclear how this impairment leads to the signs and symptoms of hereditary multiple osteochondromas.

Trichorhinophalangeal syndrome type II

The *EXT1* gene is located in a region of chromosome 8 that is deleted in people with trichorhinophalangeal syndrome type II (TRPS II). TRPS II is a condition that causes bone and joint malformations including multiple osteochondromas (described above); distinctive facial features; intellectual disability; and abnormalities of the skin, hair, teeth, sweat glands, and nails. As a result of this deletion, affected individuals are missing one copy of the *EXT1* gene in each cell. A shortage of exostosin-1 protein causes the osteochondromas in people with TRPS II. The deletion of other genes near the *EXT1* gene likely contributes to the additional features of this condition.

Other Names for This Gene

- exostoses (multiple) 1
- exostosin 1
- EXT
- EXT1_HUMAN
- Glucuronosyl-N-acetylglucosaminyl-proteoglycan 4-alpha-N-acetylglucosaminyltransferase
- N-acetylglucosaminyl-proteoglycan 4-beta-glucuronosyltransferase

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of EXT1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2131\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2131[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28EXT1%5BTIAB%5D%29+OR+%28exostoses+++1%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29%29%29>)

Catalog of Genes and Diseases from OMIM

- EXOSTOSIN GLYCOSYLTRANSFERASE 1; EXT1 (<https://omim.org/entry/608177>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/2131>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=EXT1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=EXT1[gene]))

References

- Francannet C, Cohen-Tanugi A, Le Merrer M, Munnich A, Bonaventure J, Legeai-Mallet L. Genotype-phenotype correlation in hereditary multiple exostoses. *J Med Genet*. 2001 Jul;38(7):430-4. doi: 10.1136/jmg.38.7.430. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11432960>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1757186/>)
- Lonie L, Porter DE, Fraser M, Cole T, Wise C, Yates L, Wakeling E, Blair E, Morava E, Monaco AP, Ragoussis J. Determination of the mutation spectrum of the EXT1/

EXT2 genes in British Caucasian patients with multiple osteochondromas, and exclusion of six candidate genes in EXT negative cases. Hum Mutat. 2006 Nov;27(11):1160. doi: 10.1002/humu.9467. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17041877>)

- Maas SM, Shaw AC, Bikker H, Ludecke HJ, van der Tuin K, Badura-Stronka M, Belligni E, Biamino E, Bonati MT, Carvalho DR, Cobben J, de Man SA, Den Hollander NS, Di Donato N, Garavelli L, Gronborg S, Herkert JC, Hoogeboom AJ, Jamsheer A, Latos-Bielenska A, Maat-Kievit A, Magnani C, Marcelis C, Mathijssen IB, Nielsen M, Otten E, Ousager LB, Pilch J, Plomp A, Poke G, Poluha A, Posmyk R, Rieubland C, Silengo M, Simon M, Steichen E, Stumpel C, Szakszon K, Polonkai E, van den Ende J, van der Steen A, van Essen T, van Haeringen A, van Hagen JM, Verheij JB, Mannens MM, Hennekam RC. Phenotype and genotype in 103 patients with tricho-rhino-phalangeal syndrome. Eur J Med Genet. 2015 May;58(5):279-92. doi:10.1016/j.ejmg.2015.03.002. Epub 2015 Mar 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25792522>)
- McCormick C, Duncan G, Goutsos KT, Tufaro F. The putative tumor suppressors EXT1 and EXT2 form a stable complex that accumulates in the Golgi apparatus and catalyzes the synthesis of heparan sulfate. Proc Natl Acad Sci U S A. 2000 Jan 18;97(2):668-73. doi: 10.1073/pnas.97.2.668. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10639137>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC15388/>)
- Tuysuz B, Gunes N, Alkaya DU. Trichorhinophalangeal Syndrome. 2017 Apr 20 [updated 2024 Mar 21]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. GeneReviews(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK425926/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/28426188>)
- Wuyts W, Van Hul W, De Boulle K, Hendrickx J, Bakker E, Vanhoenacker F, Mollica F, Ludecke HJ, Sayli BS, Pazzaglia UE, Mortier G, Hamel B, Conrad EU, Matsushita M, Raskind WH, Willems PJ. Mutations in the EXT1 and EXT2 genes in hereditary multiple exostoses. Am J Hum Genet. 1998 Feb;62(2):346-54. doi:10.1086/301726. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9463333>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1376901/>)

Genomic Location

The *EXT1* gene is found on chromosome 8 (<https://medlineplus.gov/genetics/chromosome/8/>).

Last updated June 1, 2017