

EYA1 gene

EYA transcriptional coactivator and phosphatase 1

Normal Function

The *EYA1* gene provides instructions for making a protein that plays a role in regulating the activity of other genes. Based on this role, the EYA1 protein is called a transcription factor or transcription coactivator.

The EYA1 protein interacts with several other proteins, including a group known as SIX proteins, to turn on (activate) and turn off (inactivate) genes that are important for normal development. Before birth, these protein interactions appear to be essential for the normal formation of many tissues. These include the second branchial arch, which gives rise to tissues in the front and side of the neck, and the eyes, ears, and kidneys. After birth, these interactions are important for normal organ function.

Health Conditions Related to Genetic Changes

Branchiootorenal/branchiootic syndrome

At least 160 mutations in the *EYA1* gene have been identified in people with branchiootorenal (BOR) syndrome, a condition that disrupts the development of tissues in the neck and causes malformations of the ears and kidneys. *EYA1* gene mutations have also been found to cause branchiootic (BO) syndrome, which includes many of the same features as BOR syndrome except for kidney (renal) malformations. The two conditions are otherwise so similar that researchers often consider them together (BOR/BO syndrome or branchiootorenal spectrum disorders).

Many of the mutations that cause BOR/BO syndrome change the 3-dimensional structure of the EYA1 protein, which prevents it from interacting effectively with other proteins. Because these protein interactions are necessary for the activation of certain genes during embryonic development, the altered EYA1 protein impairs the normal development of many tissues before birth. The major signs and symptoms of BOR/BO syndrome result from abnormal development of the second branchial arch, the ears, and (in BOR syndrome) the kidneys.

In some cases, the same *EYA1* gene mutation causes BOR syndrome in some members of a family and BO syndrome in others. This variability might result from changes in other, unidentified genes that affect how the EYA1 protein functions in the

kidneys.

Congenital anomalies of kidney and urinary tract

MedlinePlus Genetics provides information about Congenital anomalies of kidney and urinary tract

Other disorders

Several mutations in the *EYA1* gene have been associated with eye abnormalities including clouding of the lens (cataracts) and clouding of the clear front surface of the eye (the cornea). These abnormalities occur without the characteristic features of BOR/BO syndrome. Researchers believe that the *EYA1* gene mutations responsible for eye abnormalities have less severe effects on protein function than the mutations that underlie BOR/BO syndrome.

Other Names for This Gene

- BOP
- BOR
- EYA1_HUMAN
- eyes absent 1
- eyes absent homolog 1 (Drosophila)
- eyes absent, Drosophila, homolog of, 1

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of EYA1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2138\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2138[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28EYA1%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- EYA TRANSCRIPTIONAL COACTIVATOR AND PHOSPHATASE 1; EYA1 (<https://omim.org/entry/601653>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/2138>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=EYA1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=EYA1[gene]))

References

- Abdelhak S, Kalatzis V, Heilig R, Compain S, Samson D, Vincent C, Weil D, Cruaud C, Sahly I, Leibovici M, Bitner-Glindzicz M, Francis M, Lacombe D, Vigneron J, Charachon R, Boven K, Bedbeder P, Van Regemorter N, Weissenbach J, Petit C. A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal (BOR) syndrome and identifies a novel gene family. *Nat Genet.* 1997 Feb;15(2):157-64. doi: 10.1038/ng0297-157. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9020840>)
- Azuma N, Hirakiyama A, Inoue T, Asaka A, Yamada M. Mutations of a human homologue of the *Drosophila* eyes absent gene (*EYA1*) detected in patients with congenital cataracts and ocular anterior segment anomalies. *Hum Mol Genet.* 2000 Feb 12;9(3):363-6. doi: 10.1093/hmg/9.3.363. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10655545>)
- Chang EH, Menezes M, Meyer NC, Cucci RA, Vervoort VS, Schwartz CE, Smith RJ. Branchio-oto-renal syndrome: the mutation spectrum in *EYA1* and its phenotypic consequences. *Hum Mutat.* 2004 Jun;23(6):582-9. doi: 10.1002/humu.20048. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15146463>)
- Krug P, Moriniere V, Marlin S, Koubi V, Gabriel HD, Colin E, Bonneau D, Salomon R, Antignac C, Heidet L. Mutation screening of the *EYA1*, *SIX1*, and *SIX5* genes in a large cohort of patients harboring branchio-oto-renal syndrome calls into question the pathogenic role of *SIX5* mutations. *Hum Mutat.* 2011 Feb;32(2):183-90. doi: 10.1002/humu.21402. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21280147>)
- Orten DJ, Fischer SM, Sorensen JL, Radhakrishna U, Cremers CW, Marres HA, VanCamp G, Welch KO, Smith RJ, Kimberling WJ. Branchio-oto-renal syndrome (BOR): novel mutations in the *EYA1* gene, and a review of the mutational genetics of BOR. *Hum Mutat.* 2008 Apr;29(4):537-44. doi: 10.1002/humu.20691. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18220287>)
- Rayapureddi JP, Hegde RS. Branchio-oto-renal syndrome associated mutations in Eyes Absent 1 result in loss of phosphatase activity. *FEBS Lett.* 2006 Jul 10;580(16):3853-9. doi: 10.1016/j.febslet.2006.06.009. Epub 2006 Jun 15. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16797546>)
- Reis LM, Tyler RC, Muheisen S, Raggio V, Salviati L, Han DP, Costakos D, Yonath H, Hall S, Power P, Semina EV. Whole exome sequencing in dominant cataract identifies a new causative factor, *CRYBA2*, and a variety of novel alleles in known genes. *Hum Genet.* 2013 Jul;132(7):761-70. doi: 10.1007/s00439-013-1289-0. Epub 2013 Mar 19. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23508780>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3683360/>)

- Smith RJH. Branchiootorenal Spectrum Disorder. 1999 Mar 19 [updated 2018 Sep6] . In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. GeneReviews(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1380/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301554>)

Genomic Location

The *EYA1* gene is found on chromosome 8 (<https://medlineplus.gov/genetics/chromosome/8/>).

Last updated March 1, 2016