

## FGFR2 gene

fibroblast growth factor receptor 2

### Normal Function

The *FGFR2* gene provides instructions for making a protein called fibroblast growth factor receptor 2 (FGFR2). Fibroblast growth factor receptors are related proteins that are involved in important processes such as cell growth and division (proliferation), cell maturation (differentiation), bone development, formation of blood vessels (angiogenesis), wound healing, and embryonic development.

The FGFR2 protein spans the outer membrane surrounding cells, so that one end of the protein remains inside the cell and the other end projects from the outer surface of the cell. This positioning allows the FGFR2 protein to interact with specific growth factors outside the cell and to receive signals that help the cell respond to its environment. When growth factors attach to the FGFR2 protein, the receptor triggers a series of chemical reactions inside the cell that instruct the cell to undergo certain changes, such as maturing to take on specialized functions.

The FGFR2 protein plays an important role in bone growth, particularly during development before birth (embryonic development). For example, this protein signals certain immature cells in the developing embryo to become bone cells and form the head, hands, feet, and other tissues. The protein is also involved in bone remodeling, a normal process in which old bone is broken down and new bone is created to replace it.

There are several slightly different versions (isoforms) of the FGFR2 protein. Specific patterns of these isoforms are found in the body's tissues, and these patterns may change throughout growth and development.

### Health Conditions Related to Genetic Changes

#### Apert syndrome

At least ten mutations in the *FGFR2* gene have been found to cause Apert syndrome. This condition causes premature closure of the bones of the skull (craniosynostosis), leading to a misshapen head, distinctive facial features, and brain abnormalities. Affected individuals often have abnormalities of the fingers and toes, hearing and vision problems, and other signs and symptoms.

More than 98 percent of cases of Apert syndrome are caused by one of two mutations

in the *FGFR2* gene. These mutations change single protein building blocks (amino acids) in the FGFR2 protein. One mutation replaces the amino acid serine with the amino acid tryptophan at protein position 252 (written as Ser252Trp). The other mutation replaces the amino acid proline with the amino acid arginine at position 253 (written as Pro253Arg). These changes are described as "gain-of-function" because they increase the activity of the protein, leading to stronger signaling, which causes cells to mature too quickly. As a result, structures in the body develop abnormally, leading to premature fusion of bones in the skull, hands, and feet and other characteristic features of Apert syndrome.

#### Beare-Stevenson cutis gyrata syndrome

At least three mutations in the *FGFR2* gene have been found to cause Beare-Stevenson cutis gyrata syndrome, a condition that causes craniosynostosis, leading to a misshapen head and distinctive facial features, and a skin abnormality called cutis gyrata. The most common mutation replaces the amino acid tyrosine with the amino acid cysteine at position 375 in the protein (written as Tyr375Cys). The *FGFR2* gene mutations that cause Beare-Stevenson cutis gyrata syndrome appear to overactivate signaling by the FGFR2 protein, which promotes the premature fusion of bones in the skull and alters skin development.

#### Crouzon syndrome

At least 60 mutations in the *FGFR2* gene can cause Crouzon syndrome, a condition that causes craniosynostosis, leading to a misshapen head and distinctive facial features. Affected individuals can have vision, hearing, and dental problems related to abnormal skull development. Most of the mutations that cause Crouzon syndrome change single DNA building blocks (nucleotides) in the *FGFR2* gene. Insertions and deletions of a small number of nucleotides are also known to cause the disorder. These mutations in the *FGFR2* gene appear to overactivate signaling by the FGFR2 protein, which promotes premature fusion of bones in the skull.

#### Jackson-Weiss syndrome

At least six mutations in the *FGFR2* gene have been found to cause Jackson-Weiss syndrome. This condition causes craniosynostosis, leading to a misshapen head and distinctive facial features. Affected individuals also have foot abnormalities. Each of the mutations associated with Jackson-Weiss syndrome changes a single amino acid in a region of the FGFR2 protein known as the IgIII domain, which is critical for receiving signals and interacting with growth factors. The mutations appear to overactivate signaling by the FGFR2 protein, which promotes premature fusion of skull bones and affects development of bones in the feet.

#### Lacrimo-auriculo-dento-digital syndrome

At least two mutations in the *FGFR2* gene have been found to cause lacrimo-auriculo-dento-digital (LADD) syndrome. The main features of LADD syndrome are abnormal tear production, malformed ears with hearing loss, decreased saliva production, small

teeth, and hand deformities. A mutation that occurs in some people with LADD syndrome replaces the amino acid alanine with the amino acid threonine at position 628 in the FGFR2 protein (written as Ala628Thr or A628T).

The *FGFR2* gene mutations that cause LADD syndrome reduce the FGFR2 protein's ability to trigger chemical signaling within cells when it binds to its growth factor. These defects in cell signaling disrupt cell maturation and development, which results in abnormal formation of glands in the eyes and mouth, the ears, and the skeleton in people with LADD syndrome.

### Pfeiffer syndrome

More than 25 mutations in the *FGFR2* gene can cause Pfeiffer syndrome, a condition that causes craniosynostosis, leading to a misshapen head and distinctive facial features. People with this condition can also have elbow, hand, or foot abnormalities. The severity of this condition varies widely among affected individuals. Several of the mutations that cause this condition change the number of cysteine amino acids in a critical region of the FGFR2 protein known as the IgIII domain. The remaining mutations affect amino acids other than cysteine or result in an FGFR2 protein that is missing one or more amino acids. These mutations appear to overactivate signaling by the FGFR2 protein, which promotes premature fusion of skull bones and affects the development of bones in the hands and feet.

### Breast cancer

MedlinePlus Genetics provides information about Breast cancer

### Cholangiocarcinoma

MedlinePlus Genetics provides information about Cholangiocarcinoma

### Epidermal nevus

MedlinePlus Genetics provides information about Epidermal nevus

### Prostate cancer

MedlinePlus Genetics provides information about Prostate cancer

### Other disorders

Mutations in the *FGFR2* gene can cause other disorders that affect bone development. *FGFR2* gene mutations have been found in individuals with conditions called bent bone dysplasia syndrome and Antley-Bixler syndrome.

Bent bone dysplasia syndrome is characterized by the underdevelopment of certain bones, bent long bones in the arms and legs, low bone mineral density (osteopenia), and distinctive facial features. Signs and symptoms of Antley-Bixler syndrome typically include craniosynostosis, distinctive facial features, arm and leg malformations, and

genitourinary abnormalities.

The *FGFR2* gene mutations that cause these conditions appear to overactivate signaling by the FGFR2 protein, leading to skeletal abnormalities and other characteristic features of the conditions.

The particular condition that results from *FGFR2* gene mutations may be determined by the region of the gene in which the mutation occurs. It is likely that mutations in different regions of the *FGFR2* gene have varied effects on the protein, resulting in a range of protein activity and signaling. Changes in other genes or modifications of DNA that do not alter the DNA sequence (epigenetic changes) may also help determine the resulting signs and symptoms. The different factors lead to similar but distinct features of the many conditions associated with *FGFR2* gene mutations.

### Cancers

Alterations in the activity (expression) of the *FGFR2* gene are associated with certain cancers. The altered gene expression may enhance several cancer-promoting cell functions such as cell division (proliferation), cell movement, and the formation of new blood vessels (angiogenesis) that nourish a growing tumor.

The *FGFR2* gene is abnormally active (overexpressed) in certain types of stomach (gastric) cancers, and this amplification is associated with a poor disease outcome. Abnormal expression of the *FGFR2* gene is also found in patients with prostate cancer. A shift in the expression of two specific FGFR2 isoforms, IIIb and IIIc, appears to correlate with prostate cancer progression. This change in expression is complex, however, and varies depending on the type of prostate tumor. More advanced tumors may show an increase in the IIIb isoform, while other prostate tumors show a decrease in IIIb but an increase in IIIc. Altered *FGFR2* gene expression is also associated with ovarian, breast, cervical, endometrial, pancreatic, lung, and head and neck cancers.

### **Other Names for This Gene**

- bacteria-expressed kinase
- BEK
- BEK fibroblast growth factor receptor
- BEK protein tyrosine kinase
- BFR-1
- CD332
- CEK3
- CFD1
- ECT1
- FGF receptor
- FGFR2\_HUMAN
- K-SAM

- keratinocyte growth factor receptor
- KGFR
- protein tyrosine kinase, receptor like 14
- TK14
- TK25
- tyrosylprotein kinase

## Additional Information & Resources

### Tests Listed in the Genetic Testing Registry

- Tests of FGFR2 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2263\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2263[geneid]))

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28FGFR2%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D%29>)

### Catalog of Genes and Diseases from OMIM

- FIBROBLAST GROWTH FACTOR RECEPTOR 2; FGFR2 (<https://omim.org/entry/176943>)

### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/2263>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=FGFR2\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=FGFR2[gene]))

## References

- Azoury SC, Reddy S, Shukla V, Deng CX. Fibroblast Growth Factor Receptor 2( FGFR2) Mutation Related Syndromic Craniosynostosis. Int J Biol Sci.2017 Nov 2;13(12):1479-1488. doi: 10.7150/ijbs.22373. eCollection 2017. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/29230096>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5723914/>)
- Carinci F, Pezzetti F, Locci P, Becchetti E, Carls F, Avantaggiato A, Becchetti A, Carinci P, Baroni T, Bodo M. Apert and Crouzon syndromes: clinical findings, genes and extracellular matrix. J Craniofac Surg. 2005 May;16(3):361-8. doi: 10.1097/01.scs.0000157078.53871.11. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15915098>)

- Chokdeemboon C, Mahatumarat C, Rojvachiranonda N, Tongkobpetch S, Suphapeetiporn K, Shotelersuk V. FGFR1 and FGFR2 mutations in Pfeiffer syndrome. *J Craniofac Surg*. 2013 Jan;24(1):150-2. doi: 10.1097/SCS.0b013e3182646454. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23348274>)
- Ibrahimi OA, Chiu ES, McCarthy JG, Mohammadi M. Understanding the molecular basis of Apert syndrome. *Plast Reconstr Surg*. 2005 Jan;115(1):264-70. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15622262>)
- Katoh M. Therapeutics Targeting FGF Signaling Network in Human Diseases. *Trends Pharmacol Sci*. 2016 Dec;37(12):1081-1096. doi: 10.1016/j.tips.2016.10.003. Epub 2016 Oct 27. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/27992319>)
- Lajeunie E, Heuertz S, El Ghouzzi V, Martinovic J, Renier D, Le Merrer M, Bonaventure J. Mutation screening in patients with syndromic craniosynostoses indicates that a limited number of recurrent FGFR2 mutations accounts for severe forms of Pfeiffer syndrome. *Eur J Hum Genet*. 2006 Mar;14(3):289-98. doi:10.1038/sj.ejhg.5201558. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16418739>)
- Lew ED, Bae JH, Rohmann E, Wollnik B, Schlessinger J. Structural basis for reduced FGFR2 activity in LADD syndrome: Implications for FGFR autoinhibition and activation. *Proc Natl Acad Sci U S A*. 2007 Dec 11;104(50):19802-7. doi:10.1073/pnas.0709905104. Epub 2007 Dec 3. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18056630>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2148379/>)
- McGlaughlin KL, Witherow H, Dunaway DJ, David DJ, Anderson PJ. Spectrum of Antley-Bixler syndrome. *J Craniofac Surg*. 2010 Sep;21(5):1560-4. doi:10.1097/SCS.0b013e3181ec6afe. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20818252>)
- Naimi B, Latil A, Fournier G, Mangin P, Cussenot O, Berthon P. Down-regulation of (IIIb) and (IIIc) isoforms of fibroblast growth factor receptor 2 (FGFR2) is associated with malignant progression in human prostate. *Prostate*. 2002 Aug;52(3):245-52. doi: 10.1002/pros.10104. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12111699>)
- Rohmann E, Brunner HG, Kayserili H, Uyguner O, Nurnberg G, Lew ED, Dobbie A, Eswarakumar VP, Uzumcu A, Ulubil-Emeroglu M, Leroy JG, Li Y, Becker C, Lehnerdt K, Cremers CW, Yuksel-Apak M, Nurnberg P, Kubisch C, Schlessinger J, van Bokhoven H, Wollnik B. Mutations in different components of FGF signaling in LADD syndrome. *Nat Genet*. 2006 Apr;38(4):414-7. doi: 10.1038/ng1757. Epub 2006 Feb 26. Erratum In: *Nat Genet*. 2006 Apr;38(4):495. Kubisch, Christian [corrected to Kubisch, Christian]. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16501574>)
- Shams I, Rohmann E, Eswarakumar VP, Lew ED, Yuzawa S, Wollnik B, Schlessinger J, Lax I. Lacrimo-auriculo-dento-digital syndrome is caused by reduced activity of the fibroblast growth factor 10 (FGF10)-FGF receptor 2 signaling pathway. *Mol Cell Biol*. 2007 Oct;27(19):6903-12. doi: 10.1128/MCB.00544-07. Epub 2007 Aug 6. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17682060>) or Free

article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2099236/>)

- Shin EY, Lee BH, Yang JH, Shin KS, Lee GK, Yun HY, Song YJ, Park SC, Kim EG. Up-regulation and co-expression of fibroblast growth factor receptors in humangastric cancer. J Cancer Res Clin Oncol. 2000 Sep;126(9):519-28. doi:10.1007/s004320000128. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11003564>)
- Wilkie AO, Patey SJ, Kan SH, van den Ouweland AM, Hamel BC. FGFR3, its receptors, and human limb malformations: clinical and molecular correlations. Am J Med Genet. 2002 Oct 15;112(3):266-78. doi: 10.1002/ajmg.10775. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12357470>)

## Genomic Location

The *FGFR2* gene is found on chromosome 10 (<https://medlineplus.gov/genetics/chromosome/10/>).

**Last updated June 1, 2020**