

FOXC1 gene

forkhead box C1

Normal Function

The *FOXC1* gene provides instructions for making a protein that attaches (binds) to specific regions of DNA and regulates the activity of other genes. On the basis of this action, the FOXC1 protein is called a transcription factor.

The FOXC1 protein plays a critical role in early development, particularly in the formation of structures in the front part of the eye (the anterior segment). These structures include the colored part of the eye (the iris), the lens of the eye, and the clear front covering of the eye (the cornea). Studies suggest that the FOXC1 protein may also have functions in the adult eye, such as helping cells respond to oxidative stress. Oxidative stress occurs when unstable molecules called free radicals accumulate to levels that can damage or kill cells.

The FOXC1 protein is also involved in the normal development of other parts of the body, including the heart, kidneys, and brain.

Health Conditions Related to Genetic Changes

Axenfeld-Rieger syndrome

More than 50 mutations in the *FOXC1* gene have been found to cause Axenfeld-Rieger syndrome type 3, a condition that primarily affects the development of the anterior segment of the eye but can also affect other parts of the body. Many *FOXC1* gene mutations reduce the amount of functional FOXC1 protein that is produced or result in a defective protein that cannot regulate the activity of other genes. Other genetic changes (such as a duplication of the *FOXC1* gene) likely increase the amount or function of the FOXC1 protein. Having either too little or too much activity of this protein disrupts the regulation of other genes needed for normal development.

Changes in the amount or function of the FOXC1 protein impairs the development of the anterior segment of the eye, leading to the eye abnormalities characteristic of Axenfeld-Rieger syndrome. In some cases, changes involving the FOXC1 protein also cause problems with development of other parts of the body.

Peters anomaly

At least two mutations in the *FOXC1* gene have been found to cause Peters anomaly. This condition is characterized by abnormal development of the anterior segment and clouding of the cornea. The mutations that cause Peters anomaly likely disrupt the protein's ability to regulate the expression of developmental genes, especially affecting the eye. Abnormal formation of the iris, cornea, and other structures of the anterior segment leads to the features of Peters anomaly. The *FOXC1* gene mutations that cause Peters anomaly can cause other related eye disorders, such as Axenfeld-Rieger syndrome (described above), in members of the same family.

Dandy-Walker malformation

MedlinePlus Genetics provides information about Dandy-Walker malformation

Other disorders

Mutations in the *FOXC1* gene have also been identified in another eye disorder called iridogoniodysgenesis type 1. Like Axenfeld-Rieger syndrome and Peters anomaly, this condition primarily involves the anterior segment of the eye. Iridogoniodysgenesis type 1 is associated with underdevelopment of the iris and an elevated risk of increased pressure in the eye (glaucoma).

FOXC1 gene mutations have been reported in a few people with abnormalities of brain development. Mutations that change single protein building blocks (amino acids) in the *FOXC1* protein have been associated with defects of the cerebellum, which is the part of the brain that is involved in coordinating movement. Additionally, deletions of genetic material from a region of chromosome 6 that includes the *FOXC1* gene and several neighboring genes have been associated with a structural abnormality of the cerebellum known as Dandy-Walker malformation.

Other Names for This Gene

- FKHL7
- forkhead box protein C1
- forkhead, drosophila, homolog-like 7
- forkhead-related activator 3
- forkhead-related protein FKHL7
- forkhead-related transcription factor 3
- forkhead/winged helix-like transcription factor 7
- FOXC1_HUMAN
- FREAC-3
- FREAC3
- myeloid factor-delta

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of FOXC1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2296\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=2296[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28FOXC1%5BTIAB%5D%29+OR+%28forkhead+box+C1%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29%29%29>)

Catalog of Genes and Diseases from OMIM

- FORKHEAD BOX C1; FOXC1 (<https://omim.org/entry/601090>)
- ANTERIOR SEGMENT DYSGENESIS 3; ASGD3 (<https://omim.org/entry/601631>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/2296>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=FOXC1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=FOXC1[gene]))

References

- Aldinger KA, Lehmann OJ, Hudgins L, Chizhikov VV, Bassuk AG, Ades LC, Krantz ID, Dobyns WB, Millen KJ. FOXC1 is required for normal cerebellar development and is a major contributor to chromosome 6p25.3 Dandy-Walker malformation. *Nat Genet.* 2009 Sep;41(9):1037-42. doi: 10.1038/ng.422. Epub 2009 Aug 9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19668217>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2843139/>)
- Berry FB, Lines MA, Oas JM, Footz T, Underhill DA, Gage PJ, Walter MA. Functional interactions between FOXC1 and PITX2 underlie the sensitivity to FOXC1 gene dose in Axenfeld-Rieger syndrome and anterior segment dysgenesis. *Hum Mol Genet.* 2006 Mar 15;15(6):905-19. doi: 10.1093/hmg/ddl008. Epub 2006 Jan 31. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16449236>)
- Dhaene B, Meire F, Claerhout I, Kroes HY, Plomp A, Arens YH, de Ravel T, Casteels I, De Jaegere S, Hooghe S, Wuyts W, van den Ende J, Roulez F, Veenstra-Knol HE, Oldenburg RA, Giltay J, Verheij JB, de Faber JT, Menten B, DePaepe A, Kestelyn P, Leroy BP, De Baere E. Expanding the spectrum of FOXC1 and PITX2 mutations and copy number changes in patients with anterior segment malformations. *Invest Ophthalmol Vis Sci.* 2011 Jan 21;52(1):324-33. doi:10.1167/iovs.10-5309. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20881294>)

- Honkanen RA, Nishimura DY, Swiderski RE, Bennett SR, Hong S, Kwon YH, Stone EM, Sheffield VC, Alward WL. A family with Axenfeld-Rieger syndrome and Peters Anomaly caused by a point mutation (Phe112Ser) in the FOXC1 gene. *Am J Ophthalmol*. 2003 Mar;135(3):368-75. doi: 10.1016/s0002-9394(02)02061-5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12614756>)
- Lehmann OJ, Ebenezer ND, Ekong R, Ocaka L, Mungall AJ, Fraser S, McGill JI, Hitchings RA, Khaw PT, Sowden JC, Povey S, Walter MA, Bhattacharya SS, Jordan T. Ocular developmental abnormalities and glaucoma associated with interstitial 6p25 duplications and deletions. *Invest Ophthalmol Vis Sci*. 2002 Jun;43(6):1843-9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12036988>)
- Nishimura DY, Searby CC, Alward WL, Walton D, Craig JE, Mackey DA, Kawase K, Kanis AB, Patil SR, Stone EM, Sheffield VC. A spectrum of FOXC1 mutation suggests gene dosage as a mechanism for developmental defects of the anterior chamber of the eye. *Am J Hum Genet*. 2001 Feb;68(2):364-72. doi: 10.1086/318183. Epub 2001 Jan 18. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11170889>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1235270/>)
- Reis LM, Tyler RC, Volkmann Kloss BA, Schilter KF, Levin AV, Lowry RB, Zwijsen PJ, Stroh E, Broeckel U, Murray JC, Semina EV. PITX2 and FOXC1 spectrum of mutations in ocular syndromes. *Eur J Hum Genet*. 2012 Dec;20(12):1224-33. doi: 10.1038/ejhg.2012.80. Epub 2012 May 9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22569110>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3499749/>)
- Strungaru MH, Dinu I, Walter MA. Genotype-phenotype correlations in Axenfeld-Rieger malformation and glaucoma patients with FOXC1 and PITX2 mutations. *Invest Ophthalmol Vis Sci*. 2007 Jan;48(1):228-37. doi:10.1167/iovs.06-0472. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/17197537>)
- Tumer Z, Bach-Holm D. Axenfeld-Rieger syndrome and spectrum of PITX2 and FOXC1 mutations. *Eur J Hum Genet*. 2009 Dec;17(12):1527-39. doi: 10.1038/ejhg.2009.93. Epub 2009 Jun 10. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19513095>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2987033/>)
- Weisschuh N, Wolf C, Wissinger B, Gramer E. A novel mutation in the FOXC1 gene in a family with Axenfeld-Rieger syndrome and Peters' anomaly. *Clin Genet*. 2008 Nov;74(5):476-80. doi: 10.1111/j.1399-0004.2008.01025.x. Epub 2008 May 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18498376>)

Genomic Location

The *FOXC1* gene is found on chromosome 6 (<https://medlineplus.gov/genetics/chromosome/6/>).

Last updated January 1, 2014