

## GJB6 gene

gap junction protein beta 6

### Normal Function

The *GJB6* gene provides instructions for making a protein called gap junction beta 6, more commonly known as connexin 30. Connexin 30 is a member of the connexin protein family. Connexin proteins form channels called gap junctions that permit the transport of nutrients, charged atoms (ions), and signaling molecules between adjoining cells. The size of the gap junction and the types of particles that move through it are determined by the particular connexin proteins that make up the channel. Gap junctions made with connexin 30 transport potassium ions and certain small molecules.

Connexin 30 is found in several different tissues throughout the body, including the brain, inner ear, skin (especially the palms of the hands and soles of the feet), hair follicles, and nail beds. Because of its presence in the inner ear, researchers are interested in this protein's role in hearing. Hearing requires the conversion of sound waves to electrical nerve impulses. This conversion involves many processes, including maintenance of the proper level of potassium ions in the inner ear. Some studies indicate that gap junctions made with connexin 30 help to maintain the correct level of potassium ions.

### Health Conditions Related to Genetic Changes

#### Clouston syndrome

At least four *GJB6* gene mutations have been identified in people with a skin disorder called Clouston syndrome, which is also known as hidrotic ectodermal dysplasia 2. Characteristics of Clouston syndrome include fingernail abnormalities, hair loss, and thickened skin on the palms of the hands and soles of the feet. The *GJB6* gene mutations that cause Clouston syndrome change single protein building blocks (amino acids) in the connexin 30 protein. Although the effects of these mutations are not fully understood, they lead to abnormalities in the growth, division, and maturation of cells in the hair follicles, nails, and skin.

#### Nonsyndromic hearing loss

Researchers have identified a few *GJB6* gene mutations in individuals with nonsyndromic hearing loss, which is loss of hearing that is not associated with other

signs and symptoms. Mutations in this gene cause a form of nonsyndromic hearing loss called DFNA3. This form of hearing loss can either be present before a child learns to speak (prelingual) or begin after a child learns to speak (postlingual). The hearing loss ranges from mild to profound, becomes more severe over time, and particularly affects the ability to hear high-frequency sounds.

At least two *GJB6* gene mutations have been reported to cause DFNA3. Each of these mutations changes a single amino acid in connexin 30. The mutations are described as "dominant negative" because they lead to an abnormal version of connexin 30 that appears to block the formation of functional gap junctions. A shortage of these channels may alter the level of potassium ions in the inner ear, which would disrupt the conversion of sound waves to nerve impulses.

### **Other Names for This Gene**

- CX30
- CXB6\_HUMAN
- DFNA3
- ED2
- EDH
- gap junction protein, beta 6
- gap junction protein, beta 6, 30kDa
- HED

### **Additional Information & Resources**

#### Tests Listed in the Genetic Testing Registry

- Tests of GJB6 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=10804\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=10804[geneid]))

#### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28GJB6%5BTIAB%5D%29+OR+%28connexin+30%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>)

#### Catalog of Genes and Diseases from OMIM

- GAP JUNCTION PROTEIN, BETA-6; GJB6 (<https://omim.org/entry/604418>)

#### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/10804>)

- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=GJB6\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=GJB6[gene]))

## References

- Baris HN, Zlotogorski A, Peretz-Amit G, Doviner V, Shohat M, Reznik-Wolf H, Pras E. A novel GJB6 missense mutation in hidrotic ectodermal dysplasia 2 (Clouston syndrome) broadens its genotypic basis. *Br J Dermatol*. 2008 Dec;159(6):1373-6. doi: 10.1111/j.1365-2133.2008.08796.x. Epub 2008 Aug 19. No abstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18717672>)
- Fujimoto A, Kurban M, Nakamura M, Farooq M, Fujikawa H, Kibbi AG, Ito M, Dahdah M, Matta M, Diab H, Shimomura Y. GJB6, of which mutations underlie Clouston syndrome, is a potential direct target gene of p63. *J Dermatol Sci*. 2013 Feb;69(2):159-66. doi: 10.1016/j.jdermsci.2012.11.005. Epub 2012 Nov 16. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23219093>)
- Kibar Z, Dube MP, Powell J, McCuaig C, Hayflick SJ, Zonana J, Hovnanian A, Radhakrishna U, Antonarakis SE, Benohanian A, Sheeran AD, Stephan ML, Gosselin R, Kelsell DP, Christianson AL, Fraser FC, Der Kaloustian VM, Rouleau GA. Clouston hidrotic ectodermal dysplasia (HED): genetic homogeneity, presence of a founder effect in the French Canadian population and fine genetic mapping. *Eur J Hum Genet*. 2000 May;8(5):372-80. doi: 10.1038/sj.ejhg.5200471. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10854098>)
- Lamartine J, Munhoz Essfelder G, Kibar Z, Lanneluc I, Callouet E, Laoudj D, Lemaitre G, Hand C, Hayflick SJ, Zonana J, Antonarakis S, Radhakrishna U, Kelsell DP, Christianson AL, Pitaval A, Der Kaloustian V, Fraser C, Blanchet-Bardon C, Rouleau GA, Waksman G. Mutations in GJB6 cause hidrotic ectodermal dysplasia. *Nat Genet*. 2000 Oct;26(2):142-4. doi: 10.1038/79851. No abstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11017065>)
- Marlin S, Feldmann D, Blons H, Loundon N, Rouillon I, Albert S, Chauvin P, Garabedian EN, Couderc R, Odent S, Joannard A, Schmerber S, Delobel B, Leman J, Journel H, Catros H, Lemarechal C, Dollfus H, Eliot MM, Delaunoy JL, David A, Calais C, Drouin-Garraud V, Obstoy MF, Goizet C, Duriez F, Fellmann F, Helias J, Vigneron J, Montaut B, Matin-Coignard D, Faivre L, Baumann C, Lewin P, Petit C, Denoyelle F. GJB2 and GJB6 mutations: genotypic and phenotypic correlations in a large cohort of hearing-impaired patients. *Arch Otolaryngol Head Neck Surg*. 2005 Jun;131(6):481-7. doi: 10.1001/archotol.131.6.481. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15967879>)
- Mellerio J, Greenblatt D. Hidrotic Ectodermal Dysplasia 2. 2005 Apr 25 [updated 2020 Oct 15]. In: Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, Amemiya A, editors. *GeneReviews*(R) [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1200/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301379>)
- Nickel R, Forge A. Gap junctions and connexins in the inner ear: their roles in homeostasis and deafness. *Curr Opin Otolaryngol Head Neck Surg*. 2008 Oct;16(5):452-7. doi: 10.1097/MOO.0b013e32830e20b0. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18717672>)

ncbi.nlm.nih.gov/18797288)

- Smith RJH, Ranum PT. Nonsyndromic Hearing Loss and Deafness, DFNA3 - RETIREDCHAPTER, FOR HISTORICAL REFERENCE ONLY. 1998 Sep 28 [updated 2016 Dec 22]. In:Adam MP, Feldman J, Mirzaa GM, Pagon RA, Wallace SE, Bean LJH, Gripp KW, AmemiyaA, editors. GeneReviews(R) [Internet]. Seattle (WA): University ofWashington, Seattle; 1993-2024. Available from<http://www.ncbi.nlm.nih.gov/books/NBK1536/> Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20301708>)
- Xu J, Nicholson BJ. The role of connexins in ear and skin physiology -functional insights from disease-associated mutations. Biochim Biophys Acta. 2013Jan;1828(1):167-78. doi: 10.1016/j.bbame.2012.06.024. Epub 2012 Jul 13. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22796187>) or Free article on PubMed Central (<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3521577/>)

## Genomic Location

The *GJB6* gene is found on chromosome 13 (<https://medlineplus.gov/genetics/chromosome/13/>).

**Last updated February 1, 2016**