

LIPH gene

lipase H

Normal Function

The *LIPH* gene provides instructions for making an enzyme called lipase H. This enzyme is found in many cells and tissues, where it breaks down the molecule phosphatidic acid into lysophosphatidic acid (LPA) and free fatty acid. LPA is a ligand, which means that it attaches (binds) to certain proteins called receptors. A ligand and its receptor fit together like a key in a lock. LPA has multiple receptors and is involved in many cellular functions, such as cell growth and division (proliferation), cell movement (migration), and the self-destruction of cells (apoptosis).

One of LPA's receptors, the LPA₆ protein, regulates the proliferation and maturation (differentiation) of cells within hair follicles, which are specialized structures in the skin where hair growth occurs. These cell processes are important for the normal development of hair follicles and for hair growth; as the cells in the hair follicle divide, the hair strand (shaft) is pushed upward and extends beyond the skin, causing the hair to grow. Lipase H is also found in the outermost layer of skin (the epidermis) and glands in the skin that produce a substance that protects the skin and hair (sebaceous glands).

Health Conditions Related to Genetic Changes

Autosomal recessive hypotrichosis

More than 15 mutations in the *LIPH* gene have been found to cause autosomal recessive hypotrichosis, a condition that results in sparse hair growth (hypotrichosis) on the scalp and, less frequently, other parts of the body. Some mutations are specific to groups with Pakistani or Japanese ancestry, or in the Mari and Chuvash populations of Russia. *LIPH* gene mutations lead to the production of a lipase H enzyme with little or no function. Without functional lipase H, LPA is not produced. A lack of LPA impairs many cellular functions, including the proliferation and maturation of the cells that make up hair follicles. As a result, hair follicles are structurally abnormal and often underdeveloped. Irregular hair follicles alter the structure and growth of hair shafts, leading to fragile hair that breaks easily. A lack of lipase H function in the epidermis likely contributes to the skin problems sometimes seen in individuals with autosomal recessive hypotrichosis.

Other disorders

Mutations in the *LIPH* gene can also cause a hair condition called autosomal recessive woolly hair. People with this condition have hair that is unusually coarse, dry, fine, and tightly curled. Woolly hair typically affects only scalp hair and is present from birth. In some cases, affected individuals develop hypotrichosis as they get older. Certain *LIPH* gene mutations cause autosomal recessive woolly hair in some people and autosomal recessive hypotrichosis (described above) in others, even among members of the same family. Because of a shared genetic cause and overlapping features, it is uncertain whether these two conditions are separate disorders or part of the same disease spectrum.

Other Names for This Gene

- lipase member H
- lipase, member H
- LIPH_HUMAN
- LPD lipase-related protein
- LPDLR
- membrane-associated phosphatidic acid-selective phospholipase A1-alpha
- membrane-bound phosphatidic acid-selective phospholipase A1
- mPA-PLA1
- mPA-PLA1 alpha
- phospholipase A1 member B
- PLA1B

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of LIPH ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=200879\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=200879[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28LIPH%5BTIAB%5D%29+OR+%28lipase+H%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- LIPASE H; LIPH (<https://omim.org/entry/607365>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/200879>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=LIPH\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=LIPH[gene]))

References

- Horev L, Tosti A, Rosen I, Hershko K, Vincenzi C, Nanova K, Mali A, Potikha T, Zlotogorski A. Mutations in lipase H cause autosomal recessive hypotrichosis simplex with woolly hair. *J Am Acad Dermatol*. 2009 Nov;61(5):813-8. doi:10.1016/j.jaad.2009.04.020. Epub 2009 Sep 18. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19766349>)
- Khan S, Habib R, Mir H, Umm-e-Kalsoom, Naz G, Ayub M, Shafique S, Yamin T, Ali N, Basit S, Wasif N, Kamran-Ul-Hassan Naqvi S, Ali G, Wali A, Ansar M, Ahmad W. Mutations in the LPAR6 and LIPH genes underlie autosomal recessive hypotrichosis/woolly hair in 17 consanguineous families from Pakistan. *Clin Exp Dermatol*. 2011 Aug;36(6):652-4. doi: 10.1111/j.1365-2230.2011.04014.x. Epub 2011 Mar 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21426374>)
- Kurban M, Wajid M, Shimomura Y, Christiano AM. Mutations in LPAR6/P2RY5 and LIPH are associated with woolly hair and/or hypotrichosis. *J Eur Acad Dermatol Venereol*. 2013 May;27(5):545-9. doi: 10.1111/j.1468-3083.2012.04472.x. Epub 2012 Mar 5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22385360>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4296312/>)
- Shimomura Y, Wajid M, Petukhova L, Shapiro L, Christiano AM. Mutations in the lipase H gene underlie autosomal recessive woolly hair/hypotrichosis. *J Invest Dermatol*. 2009 Mar;129(3):622-8. doi: 10.1038/jid.2008.290. Epub 2008 Oct 2. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/18830268>)
- Shimomura Y, Wajid M, Zlotogorski A, Lee YJ, Rice RH, Christiano AM. Founder mutations in the lipase h gene in families with autosomal recessive woolly hair/hypotrichosis. *J Invest Dermatol*. 2009 Aug;129(8):1927-34. doi:10.1038/jid.2009.19. Epub 2009 Mar 5. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19262606>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2859194/>)
- Shinkuma S, Akiyama M, Inoue A, Aoki J, Natsuga K, Nomura T, Arita K, Abe R, Ito K, Nakamura H, Ujiie H, Shibaki A, Suga H, Tsunemi Y, Nishie W, Shimizu H. Prevalent LIPH founder mutations lead to loss of P2Y5 activation ability of PLA1alpha in autosomal recessive hypotrichosis. *Hum Mutat*. 2010 May;31(5):602-10. doi: 10.1002/humu.21235. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20213768>)
- Tanahashi K, Sugiura K, Takeichi T, Takama H, Shinkuma S, Shimizu H, Akiyama M. Prevalent founder mutation c.736T>A of LIPH in autosomal recessive woolly hair of Japanese leads to variable severity of hypotrichosis in adulthood. *J Eur Acad Dermatol Venereol*. 2013 Sep;27(9):1182-4. doi:10.1111/j.1468-3083.2012.04526.x. Epub 2012 Mar 26. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22452606>)

2449147)

Genomic Location

The *LIPH* gene is found on chromosome 3 (<https://medlineplus.gov/genetics/chromosome/3/>).

Last updated April 1, 2013