

## NTRK1 gene

neurotrophic receptor tyrosine kinase 1

### Normal Function

The *NTRK1* gene provides instructions for making a protein that is essential for the development and survival of nerve cells (neurons), especially those that transmit information about sensations such as pain, temperature, and touch (sensory neurons). The NTRK1 protein is found on the surface of cells, particularly sensory neurons. It acts as a kinase, which is an enzyme that changes the activity of other proteins by adding a cluster of oxygen and phosphorus atoms (a phosphate group) at specific positions. This process is called phosphorylation. The NTRK1 protein is turned on (activated) when another protein called nerve growth factor beta (NGF $\beta$ ) attaches (binds) to it and signals the NTRK1 protein to phosphorylate itself (autophosphorylation). Then, the activated NTRK1 protein phosphorylates other proteins; this process is needed to transmit signals for cell growth and survival.

### Health Conditions Related to Genetic Changes

#### Congenital insensitivity to pain with anhidrosis

Mutations in the *NTRK1* gene cause congenital insensitivity to pain with anhidrosis (CIPA), a condition characterized by the inability to feel pain and decreased or absent sweating (anhidrosis). Many mutations in the *NTRK1* gene are known to cause the condition. Many of the *NTRK1* gene mutations lead to a protein that cannot be activated by phosphorylation, which means the mutated NTRK1 protein cannot transmit cell growth and survival signals to neurons. Without the proper signaling, neurons die by a process of self-destruction called apoptosis. Loss of sensory neurons leads to the inability to feel pain in people with CIPA. In addition, people with CIPA lose the nerves leading to their sweat glands, which causes the anhidrosis seen in affected individuals.

#### Cancers

Mutations in the *NTRK1* gene are frequently found in people with a common type of thyroid cancer called papillary thyroid carcinoma. These mutations are acquired during a person's lifetime and are present only in certain cells. Such mutations are called somatic mutations. The mutations involved in papillary thyroid carcinoma occur when rearrangements of genetic material combine part of the *NTRK1* gene with another gene. At least three other genes are known to be involved in these rearrangements: the *TPM3*

gene, the *TPR* gene, and the *TFG* gene. All of these genetic rearrangements create mutated proteins called TRK oncoproteins. Unlike normal NTRK1 protein, TRK oncoproteins do not have to be activated by binding to the NGF $\beta$  protein; they are always turned on. Constant activation of the protein signals for the cells to grow and divide continuously, which can lead to papillary thyroid carcinoma.

### Other Names for This Gene

- high affinity nerve growth factor receptor
- MTC
- neurotrophic tyrosine kinase, receptor, type 1
- NTRK1\_HUMAN
- p140-TrkA
- TRK
- Trk-A
- TRK1
- TRK1-transforming tyrosine kinase protein
- TRKA
- tyrosine kinase receptor A

### Additional Information & Resources

#### Tests Listed in the Genetic Testing Registry

- Tests of NTRK1 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=4914\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=4914[geneid]))

#### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28NTRK1%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

#### Catalog of Genes and Diseases from OMIM

- NEUROTROPHIC TYROSINE KINASE, RECEPTOR, TYPE 1; NTRK1 (<https://omim.org/entry/191315>)
- THYROID CANCER, NONMEDULLARY, 1; NMTC1 (<https://omim.org/entry/188550>)
- TRAFFICKING FROM ER TO GOLGI REGULATOR; TFG (<https://omim.org/entry/602498>)

## Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/4914>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=NTRK1\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=NTRK1[gene]))

## References

- Greco A, Miranda C, Pierotti MA. Rearrangements of NTRK1 gene in papillary thyroid carcinoma. *Mol Cell Endocrinol*. 2010 May 28;321(1):44-9. doi:10.1016/j.mce.2009.10.009. Epub 2009 Oct 31. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19883730>)
- Indo Y, Tsuruta M, Hayashida Y, Karim MA, Ohta K, Kawano T, Mitsubuchi H, Tonoki H, Awaya Y, Matsuda I. Mutations in the TRKA/NGF receptor gene in patients with congenital insensitivity to pain with anhidrosis. *Nat Genet*. 1996 Aug;13(4):485-8. doi: 10.1038/ng0896-485. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8696348>)
- Indo Y. Molecular basis of congenital insensitivity to pain with anhidrosis (CIPA): mutations and polymorphisms in TRKA (NTRK1) gene encoding the receptor tyrosine kinase for nerve growth factor. *Hum Mutat*. 2001 Dec;18(6):462-71. doi:10.1002/humu.1224. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11748840>)
- Kaplan DR, Miller FD. Neurotrophin signal transduction in the nervous system. *Curr Opin Neurobiol*. 2000 Jun;10(3):381-91. doi: 10.1016/s0959-4388(00)00092-1. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10851172>)
- Miranda C, Di Virgilio M, Selleri S, Zanotti G, Pagliardini S, Pierotti MA, Greco A. Novel pathogenic mechanisms of congenital insensitivity to pain with anhidrosis genetic disorder unveiled by functional analysis of neurotrophin receptor tyrosine kinase type 1/nerve growth factor receptor mutations. *J Biol Chem*. 2002 Feb 22;277(8):6455-62. doi: 10.1074/jbc.M110016200. Epub 2001 Nov 21. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11719521>)
- Pierotti MA, Greco A. Oncogenic rearrangements of the NTRK1/NGF receptor. *Cancer Lett*. 2006 Jan 28;232(1):90-8. doi: 10.1016/j.canlet.2005.07.043. Epub 2005 Oct 20. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16242838>)
- Verpoorten N, De Jonghe P, Timmerman V. Disease mechanisms in hereditary sensory and autonomic neuropathies. *Neurobiol Dis*. 2006 Feb;21(2):247-55. doi:10.1016/j.nbd.2005.08.004. Epub 2005 Sep 23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16183296>)

## Genomic Location

The *NTRK1* gene is found on chromosome 1 (<https://medlineplus.gov/genetics/chromosome/1/>).

**Last updated May 1, 2011**