

## PAX8 gene

paired box 8

### Normal Function

The *PAX8* gene belongs to a family of genes that play critical roles in the formation of tissues and organs during embryonic development. The PAX gene family is also important for maintaining the normal function of certain cells after birth. To carry out these roles, the PAX genes provide instructions for making proteins that attach to specific areas of DNA. By attaching to critical DNA regions, these proteins help control the activity of particular genes (gene expression). On the basis of this action, PAX proteins are called transcription factors.

During embryonic development, the PAX8 protein is thought to activate genes involved in the formation of the kidney and the thyroid gland. The thyroid gland is a butterfly-shaped tissue in the lower neck. It releases hormones that play an important role in regulating growth, brain development, and the rate of chemical reactions in the body (metabolism). Following birth, the PAX8 protein regulates several genes involved in the production of thyroid hormones.

### Health Conditions Related to Genetic Changes

#### Congenital hypothyroidism

At least 15 mutations in this gene cause congenital hypothyroidism, a condition characterized by abnormally low levels of thyroid hormones starting from birth. Other *PAX8* gene mutations only mildly reduce thyroid hormone levels or have no detectable effect. Sometimes, identical mutations in members of the same family have different effects.

Most mutations change one of the building blocks (amino acids) used to make the PAX8 protein. Other mutations disrupt protein production, resulting in an abnormally small version of the PAX8 protein. Nearly all *PAX8* gene mutations prevent the PAX8 protein from effectively binding to DNA. One mutation alters interactions between the PAX8 protein and other transcription factors. As a result, the PAX8 protein cannot perform its role in regulating the activity of certain genes.

The thyroid gland is unusually small in people with *PAX8* gene mutations. This finding suggests that *PAX8* gene mutations disrupt the normal growth or survival of thyroid cells

during embryonic development. As a result, the thyroid gland is reduced in size and may be unable to produce the normal amount of thyroid hormones. Because cases caused by *PAX8* gene mutations are due to a problem with development of the thyroid gland, they are classified as thyroid dysgenesis.

### Tumors

The *PAX8* gene is sometimes involved in the formation of thyroid tumors (neoplasms). In these cases, abnormal growth affects particular cells called follicular thyroid cells. Some of these growths, called follicular adenomas, are noncancerous (benign). Other tumors, known as follicular carcinomas, are cancerous (malignant). In some of these neoplasms, the *PAX8* gene on chromosome 2 is fused with the *PPARG* gene on chromosome 3. The fusion results when segments of the two chromosomes are rearranged (translocated). It remains unclear how the fused *PAX8-PPARG* gene affects the growth of follicular thyroid cells, or why some neoplasms become cancerous while others are benign. It is likely that the fused gene disrupts the normal control of cell division or triggers new cell activities that promote tumor formation.

### **Other Names for This Gene**

- paired box gene 8
- PAX8\_HUMAN

### **Additional Information & Resources**

#### Tests Listed in the Genetic Testing Registry

- Tests of PAX8 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7849\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7849[geneid]))

#### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28PAX8%5BTIAB%5D%29+OR+%28paired+box+gene+8%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D%29%29%29>)

#### Catalog of Genes and Diseases from OMIM

- PAIRED BOX GENE 8; PAX8 (<https://omim.org/entry/167415>)

#### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/7849>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=PAX8\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=PAX8[gene]))

## References

- Cheung L, Messina M, Gill A, Clarkson A, Learoyd D, Delbridge L, Wentworth J, Philips J, Clifton-Bligh R, Robinson BG. Detection of the PAX8-PPAR gamma fusion oncogene in both follicular thyroid carcinomas and adenomas. *J Clin Endocrinol Metab.* 2003 Jan;88(1):354-7. doi: 10.1210/jc.2002-021020. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12519876>)
- Grasberger H, Ringkananont U, Lefrancois P, Abramowicz M, Vassart G, Refetoff S. Thyroid transcription factor 1 rescues PAX8/p300 synergism impaired by a natural PAX8 paired domain mutation with dominant negative activity. *Mol Endocrinol.* 2005 Jul;19(7):1779-91. doi: 10.1210/me.2004-0426. Epub 2005 Feb 17. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15718293>)
- Komatsu M, Takahashi T, Takahashi I, Nakamura M, Takahashi I, Takada G. Thyroid dysgenesis caused by PAX8 mutation: the hypermutability with CpG dinucleotides at codon 31. *J Pediatr.* 2001 Oct;139(4):597-9. doi:10.1067/mpd.2001.117071. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11598612>)
- Macchia PE, Lapi P, Krude H, Pirro MT, Missero C, Chiovato L, Souabni A, Baserga M, Tassi V, Pinchera A, Fenzi G, Gruters A, Busslinger M, Di Lauro R. PAX8 mutations associated with congenital hypothyroidism caused by thyroid dysgenesis. *Nat Genet.* 1998 May;19(1):83-6. doi: 10.1038/ng0598-83. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9590296>)
- Park SM, Chatterjee VK. Genetics of congenital hypothyroidism. *J Med Genet.* 2005 May;42(5):379-89. doi: 10.1136/jmg.2004.024158. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15863666>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1736062/>)
- Pasca di Magliano M, Di Lauro R, Zannini M. Pax8 has a key role in thyroid cell differentiation. *Proc Natl Acad Sci U S A.* 2000 Nov 21;97(24):13144-9. doi:10.1073/pnas.240336397. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11069301>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC27192/>)
- Raman P, Koenig RJ. Pax-8-PPAR-gamma fusion protein in thyroid carcinoma. *Nat Rev Endocrinol.* 2014 Oct;10(10):616-23. doi: 10.1038/nrendo.2014.115. Epub 2014 Jul 29. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25069464>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4290886/>)
- Ramos HE, Carre A, Chevrier L, Szinnai G, Tron E, Cerqueira TL, Leger J, Cabrol S, Puel O, Queinnec C, De Roux N, Guillot L, Castanet M, Polak M. Extreme phenotypic variability of thyroid dysgenesis in six new cases of congenital hypothyroidism due to PAX8 gene loss-of-function mutations. *Eur J Endocrinol.* 2014 Oct;171(4):499-507. doi: 10.1530/EJE-13-1006. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25214233>)
- Tell G, Pellizzari L, Esposito G, Pucillo C, Macchia PE, Di Lauro R, Damante G. Structural defects of a Pax8 mutant that give rise to congenital hypothyroidism. *Biochem J.* 1999 Jul 1;341 ( Pt 1)(Pt 1):89-93. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10377248>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1736062/>)

[nih.gov/pmc/articles/PMC1220333/](https://pubmed.ncbi.nlm.nih.gov/pmc/articles/PMC1220333/))

- Vilain C, Rydlewski C, Duprez L, Heinrichs C, Abramowicz M, Malvaux P, Renneboog B, Parma J, Costagliola S, Vassart G. Autosomal dominant transmission of congenital thyroid hypoplasia due to loss-of-function mutation of PAX8. *J Clin Endocrinol Metab*. 2001 Jan;86(1):234-8. doi: 10.1210/jcem.86.1.7140. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11232006>)

## **Genomic Location**

The *PAX8* gene is found on chromosome 2 (<https://medlineplus.gov/genetics/chromosome/2/>).

**Last updated September 1, 2015**