

PDGFB gene

platelet derived growth factor subunit B

Normal Function

The *PDGFB* gene provides instructions for making one version (isoform) of the platelet derived growth factor (PDGF) protein. This protein is involved in many cellular processes, including cell growth and division (proliferation), maturation (differentiation), and movement. The *PDGFB* gene provides instructions for a precursor protein that must be processed to be able to perform its function. Before processing, the precursor PDGFB protein attaches (binds) to another PDGFB protein or a similar protein called the PDGFA precursor protein, forming a structure known as a dimer. Once the dimer is formed, the precursor proteins are processed by being cut at specific locations, which forms the functional (active) PDGF proteins, called PDGF-BB and PDGF-AB.

The active PDGF-BB or PDGF-AB protein binds to a PDGF receptor, which initiates cellular signaling. PDGF signaling activates many pathways important in cell proliferation, differentiation, and movement.

Health Conditions Related to Genetic Changes

Dermatofibrosarcoma protuberans

Dermatofibrosarcoma protuberans, a rare type of cancer that causes a tumor in the deep layers of skin, is characterized by a somatic mutation involving the *PDGFB* gene. Somatic mutations are not inherited, but are acquired during a person's lifetime and are present only in certain cells. Dermatofibrosarcoma protuberans is associated with a rearrangement (translocation) of genetic material between chromosomes 17 and 22. This translocation, written as t(17;22), fuses part of the *PDGFB* gene on chromosome 22 with part of another gene on chromosome 17 called *COL1A1*. The translocation is found on one or more extra chromosomes that can be either the normal linear shape or circular. The resulting combined (fusion) gene is called *COL1A1-PDGFB*.

The *COL1A1-PDGFB* fusion gene provides instructions for making a fusion protein. In the translocation, the *PDGFB* gene loses the part of its DNA that inhibits its activity, and production of the COL1A1-PDGFB fusion protein is controlled by *COL1A1* gene sequences. As a result, the gene fusion leads to the production of large amounts of the fusion protein. The COL1A1-PDGFB protein forms a dimer and is processed like the normal PDGFB precursor protein. Processing removes the COL1A1 portion, which

forms a protein that researchers believe functions like the active PDGF-BB protein. Excess PDGF-BB protein abnormally stimulates cells to proliferate and differentiate, leading to the tumor formation seen in dermatofibrosarcoma protuberans.

Primary familial brain calcification

MedlinePlus Genetics provides information about Primary familial brain calcification

Other Names for This Gene

- becaplermin
- c-sis
- FLJ12858
- PDGF subunit B
- PDGF, B chain
- PDGF-2
- PDGF2
- platelet-derived growth factor 2
- platelet-derived growth factor beta polypeptide
- platelet-derived growth factor subunit B
- platelet-derived growth factor, B chain
- SIS
- SSV

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of PDGFB ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=5155\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=5155[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28PDGFB%5BTIAB%5D%29+OR+%28platelet-derived+growth+factor+beta+polypeptide%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29%29>)

Catalog of Genes and Diseases from OMIM

- PLATELET-DERIVED GROWTH FACTOR, BETA POLYPEPTIDE; PDGFB (<https://omim.org/entry/190040>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/5155>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=PDGFB\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=PDGFB[gene]))

References

- Greco A, Fusetti L, Villa R, Sozzi G, Minoletti F, Mauri P, Pierotti MA. Transforming activity of the chimeric sequence formed by the fusion of collagen gene COL1A1 and the platelet derived growth factor b-chain gene in dermatofibrosarcoma protuberans. *Oncogene*. 1998 Sep 10;17(10):1313-9. doi:10.1038/sj.onc.1202051. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9771975>)
- Heldin CH, Ostman A, Ronnstrand L. Signal transduction via platelet-derived growth factor receptors. *Biochim Biophys Acta*. 1998 Aug 19;1378(1):F79-113. doi:10.1016/s0304-419x(98)00015-8. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9739761>)
- Shimizu A, O'Brien KP, Sjoblom T, Pietras K, Buchdunger E, Collins VP, Heldin CH, Dumanski JP, Ostman A. The dermatofibrosarcoma protuberans-associated collagen type I alpha1/platelet-derived growth factor (PDGF) B-chain fusion gene generates a transforming protein that is processed to functional PDGF-BB. *Cancer Res*. 1999 Aug 1;59(15):3719-23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10446987>)
- Simon MP, Navarro M, Roux D, Pouyssegur J. Structural and functional analysis of a chimeric protein COL1A1-PDGFB generated by the translocation (17;22)(q22;q13.1) in Dermatofibrosarcoma protuberans (DP). *Oncogene*. 2001 May 24;20(23):2965-75. doi: 10.1038/sj.onc.1204426. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11420709>)
- Simon MP, Peddeutour F, Sirvent N, Grosgeorge J, Minoletti F, Coindre JM, Terrier-Lacombe MJ, Mandahl N, Craver RD, Blin N, Sozzi G, Turc-Carel C, O'Brien KP, Kedra D, Fransson I, Guilbaud C, Dumanski JP. Deregulation of the platelet-derived growth factor B-chain gene via fusion with collagen gene COL1A1 in dermatofibrosarcoma protuberans and giant-cell fibroblastoma. *Nat Genet*. 1997 Jan; 15(1):95-8. doi: 10.1038/ng0197-95. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8988177>)
- Sirvent N, Maire G, Peddeutour F. Genetics of dermatofibrosarcoma protuberans family of tumors: from ring chromosomes to tyrosine kinase inhibitor treatment. *Genes Chromosomes Cancer*. 2003 May;37(1):1-19. doi: 10.1002/gcc.10202. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12661001>)

Genomic Location

The *PDGFB* gene is found on chromosome 22 (<https://medlineplus.gov/genetics/chrom>)

osome/22/).

Last updated September 1, 2011