

## PLEC gene

plectin

### Normal Function

The *PLEC* gene provides instructions for making a protein called plectin. This protein is produced in many different tissues in the body, including skin and muscle. Within cells, plectin interacts with several molecules that make up the cell's structural framework (the cytoskeleton). For example, plectin interacts with intermediate filaments, which form networks that provide support and strength to cells. Plectin attaches (cross-links) intermediate filaments to one another and to the cell membrane.

The exact function of plectin in different tissues is unclear. In skin cells, this protein is an essential part of structures called hemidesmosomes, which attach the network of intermediate filaments to the cell membrane. It is also a component of desmosomes, which form junctions between neighboring cells. As part of these structures, plectin plays a critical role in anchoring the outer layer of the skin (the epidermis) to underlying layers.

### Health Conditions Related to Genetic Changes

#### Epidermolysis bullosa with pyloric atresia

At least nine mutations in the *PLEC* gene can cause epidermolysis bullosa with pyloric atresia (EB-PA). In addition to skin blistering, people with EB-PA are born with a life-threatening obstruction of the digestive tract called pyloric atresia. Mutations in the *PLEC* gene account for about 15 percent of all cases of EB-PA.

The *PLEC* gene mutations responsible for EB-PA change the normal structure and function of plectin. Mutations that prevent the cell from making any functional plectin tend to cause more severe signs and symptoms, while mutations that reduce the amount of plectin or alter the protein's structure usually cause milder signs and symptoms. A shortage of functional plectin disrupts the attachment of the epidermis to underlying skin layers, making the skin less resistant to friction and minor trauma. It is less clear how mutations in the *PLEC* gene are related to pyloric atresia.

#### Congenital myasthenic syndrome

MedlinePlus Genetics provides information about Congenital myasthenic syndrome

## Epidermolysis bullosa simplex

At least one mutation in the *PLEC* gene is associated with the features of epidermolysis bullosa simplex, a condition that causes the skin to be very fragile and to blister easily. This mutation has been found in a small number of families with a form of the disorder known as the Ogna type. Researchers are uncertain whether this rare condition is actually a subtype of epidermolysis bullosa simplex or represents a separate form of epidermolysis bullosa.

The mutation responsible for the Ogna type of epidermolysis bullosa simplex changes a single protein building block (amino acid) in the plectin protein. Specifically, this genetic change replaces the amino acid arginine with the amino acid tryptophan at protein position 2000 (written as Arg2000Trp or R2000W). Studies suggest that this mutation may change the way the protein folds into a 3-dimensional shape, which could prevent it from interacting with molecules that make up the cytoskeleton. In the skin, these changes cause epidermal cells to become fragile and easily damaged. As a result, the skin is less resistant to friction and minor trauma and blisters easily.

## Limb-girdle muscular dystrophy

MedlinePlus Genetics provides information about Limb-girdle muscular dystrophy

## Other disorders

Mutations in the *PLEC* gene also cause at least one other form of epidermolysis bullosa known as epidermolysis bullosa with muscular dystrophy. In addition to skin blistering, people with this disorder experience progressive muscle weakness and wasting (atrophy) later in life. More than 20 *PLEC* gene mutations have been found to cause this form of the disorder. These mutations alter the structure, function, or production of plectin. In the skin, a reduced amount of functional plectin disrupts the attachment of the epidermis to underlying skin layers, making the skin less resistant to friction. A shortage of functional plectin in muscle tissue affects critical structures needed for muscle tensing (contraction), which leads to muscle weakness and atrophy.

## **Other Names for This Gene**

- EBS1
- EBSO
- HD1
- hemidesmosomal protein 1
- PCN
- PLEC1
- PLEC1\_HUMAN
- plectin 1
- plectin 1, intermediate filament binding protein 500kDa
- Plectin-11

- PLTN

## Additional Information & Resources

### Tests Listed in the Genetic Testing Registry

- Tests of PLEC ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=5339\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=5339[geneid]))

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28PLEC1%5BTIAB%5D%29+OR+%28plectin%5BTIAB%5D%29%29+OR+%28%28HD1%5BTIAB%5D%29+OR+%28plectin+1%5BTIAB%5D%29+OR+%28PLTN%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29>)

### Catalog of Genes and Diseases from OMIM

- PLECTIN; PLEC (<https://omim.org/entry/601282>)
- EPIDERMOLYSIS BULLOSA SIMPLEX 5B, WITH MUSCULAR DYSTROPHY; EBS5B (<https://omim.org/entry/226670>)

### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/5339>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=PLEC\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=PLEC[gene]))

## References

- Charlesworth A, Chiaverini C, Chevrant-Breton J, DelRio M, Diociaiuti A, Dupuis RP, El Hachem M, Le Fiblec B, Sankari-Ho AM, Valhquist A, Wierzbicka E, Lacour JP, Meneguzzi G. Epidermolysis bullosa simplex with PLEC mutations: new phenotypes and new mutations. *Br J Dermatol*. 2013 Apr;168(4):808-14. doi:10.1111/bjd.12202. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23289980>)
- Kiritsi D, Pigors M, Tantcheva-Poor I, Wessel C, Arin MJ, Kohlhasse J, Bruckner-Tuderman L, Has C. Epidermolysis bullosa simplex Ogna revisited. *J Invest Dermatol*. 2013 Jan;133(1):270-3. doi: 10.1038/jid.2012.248. Epub 2012 Aug 2. No abstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/22854623>)
- Koss-Harnes D, Hoyheim B, Anton-Lamprecht I, Gjesti A, Jorgensen RS, Jahnsen FL, Olaisen B, Wiche G, Gedde-Dahl T Jr. A site-specific plectin mutation causes dominant epidermolysis bullosa simplex Ogna: two identical de novo

mutations. *J Invest Dermatol*. 2002 Jan;118(1):87-93. doi: 10.1046/j.0022-202x.2001.01591.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11851880>)

- Koss-Harnes D, Hoyheim B, Jonkman MF, de Groot WP, de Weerd CJ, Nikolic B, Wiche G, Gedde-Dahl T Jr. Life-long course and molecular characterization of the original Dutch family with epidermolysis bullosa simplex with muscular dystrophy due to a homozygous novel plectin point mutation. *Acta Derm Venereol*. 2004;84(2):124-31. doi: 10.1080/00015550310007094. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15206692>)
- McMillan JR, Akiyama M, Rouan F, Mellerio JE, Lane EB, Leigh IM, Owaribe K, Wiche G, Fujii N, Uitto J, Eady RA, Shimizu H. Plectin defects in epidermolysis bullosa simplex with muscular dystrophy. *Muscle Nerve*. 2007 Jan;35(1):24-35. doi:10.1002/mus.20655. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16967486>)
- Nakamura H, Sawamura D, Goto M, Nakamura H, McMillan JR, Park S, Kono S, Hasegawa S, Paku S, Nakamura T, Ogiso Y, Shimizu H. Epidermolysis bullosa simplex associated with pyloric atresia is a novel clinical subtype caused by mutations in the plectin gene (PLEC1). *J Mol Diagn*. 2005 Feb;7(1):28-35. doi:10.1016/S1525-1578(10)60005-0. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15681471>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1867514/>)
- Pfendner E, Rouan F, Uitto J. Progress in epidermolysis bullosa: the phenotypic spectrum of plectin mutations. *Exp Dermatol*. 2005 Apr;14(4):241-9. doi: 10.1111/j.0906-6705.2005.00324.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15810881>)
- Rezniczek GA, Walko G, Wiche G. Plectin gene defects lead to various forms of epidermolysis bullosa simplex. *Dermatol Clin*. 2010 Jan;28(1):33-41. doi:10.1016/j.det.2009.10.004. Erratum In: *Dermatol Clin*. 2010 Apr;28(2):439-41. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19945614>)
- Uitto J, Pulkkinen L, Smith FJ, McLean WH. Plectin and human genetic disorders of the skin and muscle. The paradigm of epidermolysis bullosa with muscular dystrophy. *Exp Dermatol*. 1996 Oct;5(5):237-46. doi:10.1111/j.1600-0625.1996.tb00124.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/8981021>)

## Genomic Location

The *PLEC* gene is found on chromosome 8 (<https://medlineplus.gov/genetics/chromosome/8/>).

**Last updated May 1, 2013**