

SDHAF2 gene

succinate dehydrogenase complex assembly factor 2

Normal Function

The *SDHAF2* gene provides instructions for making a protein that interacts with the succinate dehydrogenase (SDH) enzyme. The SDHAF2 protein helps a molecule called FAD attach to the SDH enzyme. FAD is called a cofactor because it helps the enzyme carry out its function. The FAD cofactor is required for SDH enzyme activity.

The SDH enzyme plays a critical role in mitochondria, which are structures inside cells that convert the energy from food into a form that cells can use. Within mitochondria, the SDH enzyme links two important pathways in energy conversion: the citric acid cycle (or Krebs cycle) and oxidative phosphorylation. As part of the citric acid cycle, the SDH enzyme converts a compound called succinate to another compound called fumarate.

Succinate, the compound on which the SDH enzyme acts, is an oxygen sensor in the cell and can help turn on specific pathways that stimulate cells to grow in a low-oxygen environment (hypoxia). In particular, succinate stabilizes a protein called hypoxia-inducible factor (HIF) by preventing a reaction that would allow HIF to be broken down. HIF controls several important genes involved in cell division and the formation of new blood vessels in a hypoxic environment.

The *SDHAF2* gene is a tumor suppressor, which means it prevents cells from growing and dividing in an uncontrolled way.

Health Conditions Related to Genetic Changes

Hereditary paraganglioma-pheochromocytoma

At least one mutation in the *SDHAF2* gene has been identified in people with hereditary paraganglioma-pheochromocytoma type 2. People with this condition have paragangliomas, pheochromocytomas, or both. These noncancerous (benign) tumors are associated with the nervous system. The mutation replaces a protein building block (amino acid) in the SDHAF2 protein. Specifically, the amino acid glycine is replaced with the amino acid arginine at position 78 (written as Gly78Arg or G78R). The interaction between the mutated SDHAF2 protein and the SDH complex is impaired, and attachment of the FAD cofactor is decreased. As a result, the SDH enzyme is

nonfunctional. Because the mutated SDH enzyme cannot convert succinate to fumarate, succinate accumulates in the cell. Excess succinate abnormally stabilizes HIF, which also builds up in cells. Excess HIF stimulates cells to divide and triggers the production of blood vessels when they are not needed. Rapid and uncontrolled cell division, along with the formation of new blood vessels, can lead to the development of tumors in people with hereditary paraganglioma-pheochromocytoma.

Other Names for This Gene

- C11orf79
- FLJ20487
- hSDH5
- PGL2
- SDH assembly factor 2
- SDH5
- SDHF2_HUMAN
- succinate dehydrogenase assembly factor 2, mitochondrial
- succinate dehydrogenase subunit 5, mitochondrial

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of SDHAF2 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=54949\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=54949[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28SDHAF2%5BTIAB%5D%29+OR+%28%28PGL2%5BTIAB%5D%29+OR+%28SDH5%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2160+days%22%5Bdp%5D%29>)

Catalog of Genes and Diseases from OMIM

- SUCCINATE DEHYDROGENASE COMPLEX ASSEMBLY FACTOR 2; SDHAF2 (<https://omim.org/entry/613019>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/54949>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=SDHAF2\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=SDHAF2[gene]))

References

- Bayley JP, Kunst HP, Cascon A, Sampietro ML, Gaal J, Korpershoek E, Hinojar-Gutierrez A, Timmers HJ, Hoefsloot LH, Hermsen MA, Suarez C, Hussain AK, Vriends AH, Hes FJ, Jansen JC, Tops CM, Corssmit EP, de Krijff P, Lenders JW, Cremers CW, Devilee P, Dinjens WN, de Krijger RR, Robledo M. SDHAF2 mutations in familial and sporadic paraganglioma and pheochromocytoma. *Lancet Oncol.* 2010 Apr;11(4):366-72. doi: 10.1016/S1470-2045(10)70007-3. Epub 2010 Jan 11. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20071235>)
- Hao HX, Khalimonchuk O, Schraders M, Dephoute N, Bayley JP, Kunst H, Devilee P, Cremers CW, Schiffman JD, Bentz BG, Gygi SP, Winge DR, Kremer H, Rutter J. SDH5, a gene required for flavination of succinate dehydrogenase, is mutated in paraganglioma. *Science.* 2009 Aug 28;325(5944):1139-42. doi:10.1126/science.1175689. Epub 2009 Jul 23. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19628817>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3881419/>)
- Muller U. Pathological mechanisms and parent-of-origin effects in hereditary paraganglioma/pheochromocytoma (PGL/PCC). *Neurogenetics.* 2011 Aug;12(3):175-81. doi: 10.1007/s10048-011-0280-y. Epub 2011 Mar 9. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/21547462>)
- Pasini B, Stratakis CA. SDH mutations in tumorigenesis and inherited endocrine tumours: lesson from the pheochromocytoma-paraganglioma syndromes. *J Intern Med.* 2009 Jul;266(1):19-42. doi: 10.1111/j.1365-2796.2009.02111.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/19522823>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3163304/>)
- Pollard PJ, Briere JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell M, Olpin S, Moat SJ, Hargreaves IP, Heales SJ, Chung YL, Griffiths JR, Dalglish A, McGrath JA, Gleeson MJ, Hodgson SV, Poulson R, Rustin P, Tomlinson IP. Accumulation of Krebs cycle intermediates and over-expression of HIF1alpha in tumours which result from germline FH and SDH mutations. *Hum Mol Genet.* 2005 Aug;14(15):2231-9. doi: 10.1093/hmg/ddi227. Epub 2005 Jun 29. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15987702>)
- Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD, Pan Y, Simon MC, Thompson CB, Gottlieb E. Succinate links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase. *Cancer Cell.* 2005 Jan;7(1):77-85. doi: 10.1016/j.ccr.2004.11.022. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/15652751>)

Genomic Location

The *SDHAF2* gene is found on chromosome 11 (<https://medlineplus.gov/genetics/chromosome/11/>).

Last updated June 1, 2011