

## TGFB3 gene

transforming growth factor beta 3

### Normal Function

The *TGFB3* gene provides instructions for producing a protein called transforming growth factor beta-3 (TGF $\beta$ -3). This protein is found throughout the body and is required for development before birth and throughout life. To carry out its functions, TGF $\beta$ -3 attaches (binds) to receptor proteins on the surface of cells. This binding triggers the transmission of signals within the cell, controlling various cellular activities. As part of a signaling pathway, called the TGF- $\beta$  pathway, the TGF $\beta$ -3 protein helps control the growth and division (proliferation) of cells, the process by which cells mature to carry out specific functions (differentiation), cell movement (motility), and controlled cell death (apoptosis). Because the TGF $\beta$ -3 protein keeps cells from growing and dividing too rapidly or in an uncontrolled way, it can suppress the formation of tumors.

The TGF $\beta$ -3 protein is especially abundant in tissues that develop into the muscles used for movement (skeletal muscles), and plays a key role in their development. The protein is also involved in the formation of blood vessels, regulation of bone growth, wound healing, and immune system function.

### Health Conditions Related to Genetic Changes

#### Loeys-Dietz syndrome

At least 11 mutations in the *TGFB3* gene have been found to cause Loeys-Dietz syndrome type V. This disorder affects connective tissue, which gives structure and support to blood vessels, the skeleton, and many other parts of the body. Loeys-Dietz syndrome type V is characterized by blood vessel abnormalities, heart defects, and skeletal deformities. The *TGFB3* gene mutations that cause this condition lead to the production of a TGF $\beta$ -3 protein with little or no function. As a result, the protein cannot bind to its receptors. Although the TGF $\beta$ -3 protein and its receptors are not bound, TGF- $\beta$  pathway signaling occurs at an even greater intensity than normal. Researchers speculate that the activity of other proteins in this signaling pathway is increased to compensate for the reduction in TGF $\beta$ -3 activity; however, the exact mechanism responsible for the increase in signaling is unclear. The overactive signaling pathway disrupts development of connective tissue and various body systems and leads to the signs and symptoms of Loeys-Dietz syndrome type V.

## Arrhythmogenic right ventricular cardiomyopathy

MedlinePlus Genetics provides information about Arrhythmogenic right ventricular cardiomyopathy

### **Other Names for This Gene**

- RNHF
- TGF beta 3
- TGF-beta3

### **Additional Information & Resources**

#### Tests Listed in the Genetic Testing Registry

- Tests of TGFB3 ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7043\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7043[geneid]))

#### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28TGFB3%5BTIAB%5D%29+OR+%28transforming+growth+factor+beta+3%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D%29>)

#### Catalog of Genes and Diseases from OMIM

- TRANSFORMING GROWTH FACTOR, BETA-3; TGFB3 (<https://omim.org/entry/190230>)

#### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/7043>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=TGFB3\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=TGFB3[gene]))

### **References**

- Bertoli-Avella AM, Gillis E, Morisaki H, Verhagen JMA, de Graaf BM, van deBeek G, Gallo E, Kruithof BPT, Venselaar H, Myers LA, Laga S, Doyle AJ, Oswald G, van Cappellen GWA, Yamanaka I, van der Helm RM, Beverloo B, de Klein A, Pardo L, Lammens M, Evers C, Devriendt K, Dumoulein M, Timmermans J, Bruggenwirth HT, Verheijen F, Rodrigus I, Baynam G, Kempers M, Saenen J, Van Craenenbroeck EM, Minatoya K, Matsukawa R, Tsukube T, Kubo N, Hofstra R, Goumans MJ, Bekkers

JA, Roos-Hesselink JW, van de Laar IMBH, Dietz HC, Van Laer L, Morisaki T, Wessels MW, Loeys BL. Mutations in a TGF-beta ligand, TGFB3, cause syndromic aortic aneurysms and dissections. *J Am Coll Cardiol*. 2015 Apr 7;65(13):1324-1336. doi:10.1016/j.jacc.2015.01.040. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/25835445>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4380321/>)

- Matyas G, Naef P, Tollens M, Oexle K. De novo mutation of the latency-associated peptide domain of TGFB3 in a patient with overgrowth and Loeys-Dietz syndrome features. *Am J Med Genet A*. 2014 Aug;164A(8):2141-3. doi:10.1002/ajmg.a.36593. Epub 2014 May 5. No abstract available. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24798638>)
- Rienhoff HY Jr, Yeo CY, Morissette R, Khrebtukova I, Melnick J, Luo S, Leng N, Kim YJ, Schroth G, Westwick J, Vogel H, McDonnell N, Hall JG, Whitman M. A mutation in TGFB3 associated with a syndrome of low muscle mass, growth retardation, distal arthrogryposis and clinical features overlapping with Marfan and Loeys-Dietz syndrome. *Am J Med Genet A*. 2013 Aug;161A(8):2040-6. doi:10.1002/ajmg.a.36056. Epub 2013 Jul 3. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23824657>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3885154/>)

## Genomic Location

The *TGFB3* gene is found on chromosome 14 (<https://medlineplus.gov/genetics/chromosome/14/>).

**Last updated March 1, 2017**