

## TGFBI gene

transforming growth factor beta induced

### Normal Function

The *TGFBI* gene provides instructions for making a protein called transforming growth factor beta induced (TGFBI). This protein is released (secreted) from cells and becomes part of the extracellular matrix, which is an intricate network that forms in the spaces between cells and provides structural support to tissues. The TGFBI protein is thought to play a role in the attachment of cells to one another (cell adhesion) and cell movement (migration). This protein is found in many tissues in the body, including the the clear, outer covering of the eye (the cornea).

### Health Conditions Related to Genetic Changes

#### Lattice corneal dystrophy type I

At least 26 mutations in the *TGFBI* gene can cause lattice corneal dystrophy type I. This inherited eye condition is characterized by a lattice-like accumulation of proteins (called amyloid deposits) that form in the cornea. These deposits cloud the cornea and lead to vision impairment in affected individuals. The cornea is made up of several layers of tissue. In lattice corneal dystrophy type I, the deposits occur in the stromal layer of the cornea and contain altered TGFBI proteins.

The *TGFBI* gene mutations involved in lattice corneal dystrophy type I change single protein building blocks (amino acids) in the TGFBI protein. The most common mutation replaces the amino acid arginine with the amino acid cysteine at position 124 (written as Arg124Cys or R124C). When the condition is caused by this mutation, it is often called classic lattice corneal dystrophy type I. Altered TGFBI proteins abnormally clump together and form amyloid deposits. However, it is unclear how the changes caused by the gene mutations induce the protein to form these deposits.

#### Keratoconus

MedlinePlus Genetics provides information about Keratoconus

#### Other disorders

*TGFBI* gene mutations are involved in a number of other types of corneal dystrophy,

including Groenouw corneal dystrophy, Avellino corneal dystrophy, lattice corneal dystrophy type IIIA, Reis-Bucklers corneal dystrophy, Thiel-Behnke corneal dystrophy, and epithelial basement membrane corneal dystrophy. Corneal dystrophies are all characterized by the accumulation of protein deposits in the cornea, which leads to vision impairment. These deposits form in various shapes and patterns and occur in different layers of the cornea, which help define the different types of corneal dystrophies.

The *TGFBI* gene mutations that cause these conditions change single amino acids in the TGFBI protein. Most mutations alter the amino acids found at positions 124 and 555 of the protein, although other amino acid changes can also occur. Certain mutations are strongly associated with particular corneal dystrophies. For instance, the mutation that replaces the amino acid arginine at position 124 with the amino acid histidine (written as Arg124His) most often causes Avellino corneal dystrophy. The mutation that replaces the amino acid arginine at position 124 with the amino acid lysine (written as Arg124Lys) most often causes Reis-Bucklers corneal dystrophy. The mutation that replaces the amino acid arginine at position 555 with the amino acid glutamine (written as Arg555Gln) most often causes Thiel-Behnke corneal dystrophy. And, the mutation that replaces the amino acid arginine at position 555 with the amino acid tryptophan (written as Arg555Trp) most often causes Groenouw corneal dystrophy.

### Other Names for This Gene

- beta ig-h3
- BGH3\_HUMAN
- BIGH3
- CDB1
- CDG2
- CDGG1
- CSD
- CSD1
- CSD2
- CSD3
- EBMD
- kerato-epithelin
- LCD1
- RGD-CAP
- RGD-containing collagen-associated protein
- transforming growth factor, beta-induced, 68kDa
- transforming growth factor-beta-induced protein ig-h3

## Additional Information & Resources

### Tests Listed in the Genetic Testing Registry

- Tests of TGFBI ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7045\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7045[geneid]))

### Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28TGFBI%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+720+days%22%5Bdp%5D>)

### Catalog of Genes and Diseases from OMIM

- CORNEAL DYSTROPHY, EPITHELIAL BASEMENT MEMBRANE; EBMD (<https://omim.org/entry/121820>)
- CORNEAL DYSTROPHY, GROENOUW TYPE I; CDGG1 (<https://omim.org/entry/121900>)
- TRANSFORMING GROWTH FACTOR, BETA-INDUCED, 68-KD; TGFBI (<https://omim.org/entry/601692>)
- CORNEAL DYSTROPHY, THIEL-BEHNKE TYPE; CDTB (<https://omim.org/entry/602082>)
- CORNEAL DYSTROPHY, REIS-BUCKLERS TYPE; CDRB (<https://omim.org/entry/608470>)
- CORNEAL DYSTROPHY, LATTICE TYPE IIIA; CDL3A (<https://omim.org/entry/608471>)
- CORNEAL DYSTROPHY, AVELLINO TYPE; CDA (<https://omim.org/entry/607541>)

### Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/7045>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=TGFBI\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=TGFBI[gene]))

## References

- Boutboul S, Black GC, Moore JE, Sinton J, Menasche M, Munier FL, Laroche L, Abitbol M, Schorderet DF. A subset of patients with epithelial basement membranecorneal dystrophy have mutations in TGFBI/BIGH3. Hum Mutat. 2006 Jun; 27(6):553-7.doi: 10.1002/humu.20331. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16652336>)
- Clout NJ, Hohenester E. A model of FAS1 domain 4 of the corneal proteinbeta(ig)-h3 gives a clearer view on corneal dystrophies. Mol Vis. 2003 Sep11;9:440-8.

Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/14502125>)

- Kannabiran C, Klintworth GK. TGFB1 gene mutations in corneal dystrophies. Hum Mutat. 2006 Jul;27(7):615-25. doi: 10.1002/humu.20334. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/16683255>)
- Munier FL, Korvatska E, Djemai A, Le Paslier D, Zografos L, Pescia G, Schorderet DF. Kerato-epithelin mutations in four 5q31-linked corneal dystrophies. Nat Genet. 1997 Mar;15(3):247-51. doi: 10.1038/ng0397-247. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/9054935>)
- Schmitt-Bernard CF, Chavanieu A, Derancourt J, Arnaud B, Demaille JG, Calas B, Argiles A. In vitro creation of amyloid fibrils from native and Arg124Cys mutated beta1GH3((110-131)) peptides, and its relevance for lattice corneal amyloid dystrophy type I. Biochem Biophys Res Commun. 2000 Jul 5;273(2):649-53. doi:10.1006/bbrc.2000.2955. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/10873659>)
- Yang J, Han X, Huang D, Yu L, Zhu Y, Tong Y, Zhu B, Li C, Weng M, Ma X. Analysis of TGFB1 gene mutations in Chinese patients with corneal dystrophies and review of the literature. Mol Vis. 2010 Jun 30;16:1186-93. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/20664689>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2901189/>)

## Genomic Location

The *TGFB1* gene is found on chromosome 5 (<https://medlineplus.gov/genetics/chromosome/5/>).

**Last updated April 1, 2012**