

TPO gene

thyroid peroxidase

Normal Function

The *TPO* gene provides instructions for making an enzyme called thyroid peroxidase. This enzyme plays a central role in the function of the thyroid gland, a butterfly-shaped tissue in the lower neck. Thyroid peroxidase assists the chemical reaction that adds iodine to a protein called thyroglobulin, a critical step in generating thyroid hormones. Thyroid hormones play an important role in regulating growth, brain development, and the rate of chemical reactions in the body (metabolism).

To function properly, thyroid peroxidase must be located in the cell membrane of certain thyroid cells, called follicular cells. Thyroid peroxidase has several different versions (isoforms), which vary by size and location within the cell. Some versions do not function because they are not located in the cell membrane.

Health Conditions Related to Genetic Changes

Congenital hypothyroidism

TPO gene mutations can cause congenital hypothyroidism, a condition characterized by abnormally low levels of thyroid hormones starting from birth. The *TPO* gene mutations involved in this condition delete, add, or change DNA building blocks (base pairs) in the *TPO* gene. Some mutations lead to an abnormally small thyroid peroxidase enzyme that breaks apart before it can be inserted into the cell membrane. Other mutations change the enzyme's 3-dimensional shape, preventing it from functioning properly within the cell membrane. Without functional thyroid peroxidase, iodine taken up by the thyroid gland is not added to thyroglobulin. As a result, the production of thyroid hormones is absent or reduced, leading to the features of congenital hypothyroidism. In most affected individuals, the thyroid gland is enlarged (goiter) in an attempt to compensate for reduced hormone production. Because cases caused by *TPO* gene mutations result from a disruption of thyroid hormone synthesis, they are classified as thyroid dysmorphogenesis.

Other Names for This Gene

- MSA
- PERT_HUMAN

- thyroid microsomal antigen
- thyroperoxidase
- TPX

Additional Information & Resources

Tests Listed in the Genetic Testing Registry

- Tests of TPO ([https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7173\[geneid\]](https://www.ncbi.nlm.nih.gov/gtr/all/tests/?term=7173[geneid]))

Scientific Articles on PubMed

- PubMed (<https://pubmed.ncbi.nlm.nih.gov/?term=%28%28TPO%5BTIAB%5D%29+OR+%28thyroid+peroxidase%5BTIAB%5D%29%29+AND+%28iodide+peroxidase%5BMAJR%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>)

Catalog of Genes and Diseases from OMIM

- THYROID PEROXIDASE; TPO (<https://omim.org/entry/606765>)

Gene and Variant Databases

- NCBI Gene (<https://www.ncbi.nlm.nih.gov/gene/7173>)
- ClinVar ([https://www.ncbi.nlm.nih.gov/clinvar?term=TPO\[gene\]](https://www.ncbi.nlm.nih.gov/clinvar?term=TPO[gene]))

References

- Bakker B, Bikker H, Vulsma T, de Randamie JS, Wiedijk BM, De Vijlder JJ. Twodecades of screening for congenital hypothyroidism in The Netherlands: TPO genemutations in total iodide organification defects (an update). J Clin EndocrinolMetab. 2000 Oct;85(10):3708-12. doi: 10.1210/jcem.85.10.6878. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/11061528>)
- Cangul H, Aycan Z, Olivera-Nappa A, Saglam H, Schoenmakers NA, Boelaert K, Cetinkaya S, Tarim O, Bober E, Darendeliler F, Bas V, Demir K, Aydin BK, KendallIM, Cole T, Hogler W, Chatterjee VK, Barrett TG, Maher ER. Thyroiddys hormonogenesis is mainly caused by TPO mutations in consanguineous community.Clin Endocrinol (Oxf). 2013 Aug;79(2):275-81. doi: 10.1111/cen.12127. Epub 2013May 6. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/23236987>)
- Chardes T, Chapal N, Bresson D, Bes C, Giudicelli V, Lefranc MP, Peraldi-RouxS. The human anti-thyroid peroxidase autoantibody repertoire in Graves'

and Hashimoto's autoimmune thyroid diseases. *Immunogenetics*. 2002 Jun;54(3):141-57. doi: 10.1007/s00251-002-0453-9. Epub 2002 May 3. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12073143>)

- Kotani T, Umeki K, Kawano J, Suganuma T, Hishinuma A, Ieiri T, Harada S. Partial iodide organification defect caused by a novel mutation of the thyroid peroxidase gene in three siblings. *Clin Endocrinol (Oxf)*. 2003 Aug;59(2):198-206. doi: 10.1046/j.1365-2265.2003.01823.x. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12864797>)
- Lee CC, Harun F, Jalaludin MY, Lim CY, Ng KL, Mat Junit S. Functional analyses of C.2268dup in thyroid peroxidase gene associated with goitrous congenital hypothyroidism. *Biomed Res Int*. 2014;2014:370538. doi: 10.1155/2014/370538. Epub 2014 Mar 17. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/24745015>) or Free article on PubMed Central (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3976875/>)
- Nascimento AC, Guedes DR, Santos CS, Knobel M, Rubio IG, Medeiros-Neto G. Thyroperoxidase gene mutations in congenital goitrous hypothyroidism with total and partial iodide organification defect. *Thyroid*. 2003 Dec;13(12):1145-51. doi:10.1089/10507250360731550. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/14751036>)
- Rivolta CM, Esperante SA, Gruneiro-Papendieck L, Chiesa A, Moya CM, Domene S, Varela V, Targovnik HM. Five novel inactivating mutations in the thyroid peroxidase gene responsible for congenital goiter and iodide organification defect. *Hum Mutat*. 2003 Sep;22(3):259. doi: 10.1002/humu.9175. Citation on PubMed (<https://pubmed.ncbi.nlm.nih.gov/12938097>)

Genomic Location

The *TPO* gene is found on chromosome 2 (<https://medlineplus.gov/genetics/chromosome/2/>).

Last updated September 1, 2015